




# RHEEDEA

Journal of the Indian Association for Angiosperm Taxonomy

Vol. 09(2): 155-158 (1999)  
ISSN: 0971-2313 (Print edition)  
ISSN: 2582-2438 (Online edition)

## A new species of *Syzygium* Gaertn. (Myrtaceae) from Southern Western Ghats, India

Sasidharan N. & Jomy Augustine



How to cite:

**Sasidharan N. & J. Augustine. 1999.** A new species of *Syzygium* Gaertn. (Myrtaceae) from Southern Western Ghats, India. *Rheedeia* 9(2): 155–158.

<https://dx.doi.org/10.22244/rheedeia.1999.09.02.04>

Published in print: 31.12.1999

Published Online: 01.01.2022



विज्ञान एवं प्रौद्योगिकी विभाग  
DEPARTMENT OF  
SCIENCE & TECHNOLOGY

**Published by Indian Association for Angiosperm Taxonomy**

This volume of *Rheedeia* is published with partial financial assistance from Department of Science and Technology, Government of India, New Delhi

## A new species of *Syzygium* Gaertn. (Myrtaceae) from Southern Western Ghats, India.

N. Sasidharan and Jomy Augustine<sup>1</sup>

Kerala Forest Research Institute, Peechi 680653, Kerala

---

### Abstract

A new species of *Syzygium* Gaertn. is described from Southern Western Ghats.

### INTRODUCTION

During our study on the flora of Periyar Tiger Reserve, Idukki district, Kerala, 18 taxa of Myrtaceae were collected. Following the studies of Schmid (1972), 14 of them belonged to *Syzygium* Gaertn. and the rest to *Eugenia* L. One of the *Syzygium* collections was found not agreeing with any of the known species from India. The specimens were further compared with the collections from the Indo-Malesian region at Kew Herbarium and was found not matching. It is described and illustrated here as a new species as *S. periyarensis*, named after the type locality, The Periyar Tiger Reserve.

*Syzygium periyarensis* Jomy & Sasidharan sp. nov.

*Syzygio firmo* Thw. affinis in foliis obovatis ad late ellipticis coriaceis basi acuto vel obtuso cymis terminalibus corymbosis, sed laminae marginibus planis venis lateralibus tenuibus nervis intramarginalibus inconspicuis floribus maioribus differt.

*Typus* : India, Kerala state, Idukki district, Periyar Tiger Reserve, Sunderamala, ± 1300m, Jomy Augustine 13570, March 14, 1994 (Holo MH, Iso KFRI, CALI)

Evergreen trees, to 15 m high, bark smooth; branchlets terete, ca 4 mm thick. Leaves opposite, lamina 11-15 x 7-9 cm, obovate or broadly elliptic, obtusely acute, acumen ca 0.5 cm long, base acute or obtuse, coriaceous, margins flat; lateral nerves 7-14 pairs, inconspicuous above and prominent below, irregular, faint towards the margins, intramarginal veins absent or visible only towards the distal half, faint, 3-4 mm from the margin, intercostae indistinct, sparingly black punctate below; petiole 7-15 mm long, stout, dark brown. Cymes corymbose, 5-8 cm across, terminal, few-flowered; pedicel 4.5 mm long; pseudo-pedicel 3 mm long; calyx tube 12 x 12 mm, tube above the ovary 3-4 mm high, lobes 4, ca 6 x 12 mm, broadly ovate, obtuse; petals 4, white, ca 13 mm across, orbicular, concave; stamens numerous, many seriate, filaments 11-18

---

<sup>1</sup> Department of Botany, St. Thomas College, Pala, Kottayam, Kerala

N. Sasidharan and Jomy Augustine



Fig. 1 A. flowering twig, B. flower, C. L.S. of flower, D. Calyx, E. petal, F<sub>1</sub> & F<sub>2</sub> anthers

***Syzygium periyarensis***

mm long, inflexed in bud; anthers 1.5 mm long, ovate, obtuse; disk prominent, 2-3 mm, thick, shortly crenate; ovary conical, cells 2.5 x 2 mm; ovules many, placentation axile, style *ca* 22 mm long; stigma indistinct.

*Syzygium periyarensis* is allied to *S. firmum* Thw. by the obovate to broadly elliptic coriaceous leaves with acute or obtuse base and terminal corymbose cymes. But can be distinguished by the flat leaf margins, faint intramarginal nerves and larger flowers against the revolute leaf margins, two tier prominent intra-marginal nerves which are more or less looped and smaller flowers of *S. firmum* Thw. (Table 1).

**Table 1. Comparison of *Syzygium periyarensis* with *Syzygium firmum***

No.	Characters	<i>S. periyarensis</i>	<i>S. firmum</i>
1.	Leaf margin	flat	prominently revolute
2.	Lateral nerves	7-14 pairs inconspicuous above, faint towards the margin	many, equally prominent on both sides, reticulate
3.	Intramarginal vein	absent or visible only towards the distal half, 3-4 mm inside the margin	distinct, 2-tiered, prominent, looped 3-6 mm inside the margin
4.	Petiole	7-15 mm long	12-25 mm long
5.	Cymes	terminal	terminal or axillary
6.	Pedicel	4 mm long	flowers subsessile
7.	Calyx tube	12 x 12 mm	10 x 7 mm
8.	Petals	13 x 13 mm	6 x 5 mm
9.	Stamens	11-18 mm long	20-22 mm long

*Flowering* : March - April

*Habitat* : Occasional along the banks of streams in association with *Syzygium hemisphericum* (Wight) Alston, *Gordonia obtusa* Wall. ex Wight & Arn., *Ternstroemia japonica* Thunb., etc.

**N. Sasidharan and Jomy Augustine**

**Acknowledgements**

We are grateful to the Wildlife Wing of Kerala Forest Department for financial assistance for the study. Sincere thanks are due to Dr. J.F. Veldkamp, Rijksherbarium, Leiden for providing the Latin diagnosis. The first author is thankful to Ms. E. Nic Laughadha, Kew Herbarium for the help rendered in referring the relevant *Syzygium* specimens.

**Literature cited**

Schmid, R. 1972. *A Resolution of the Eugenia-Syzygium Controversy* (Myrtaceae). American Journ. Botany 59 : 423-436.