


A new species of *lone* Lindl. (Orchidaceae) from Arunachal Pradesh, India

Nageswara Rao A.



How to cite:

Nageswara Rao A. 1997. A new species of *lone* Lindl. (Orchidaceae) from Arunachal Pradesh, India. *Rheedea* 7(1): 47–50.

https://dx.doi.org/10.22244/rheedea.1997.07.01.09

Published in print: 30.06.1997 Published Online: 01.01.2022



Rheedea

Vol. 7(1): 47-50. 1997

A new species of *Ione* Lindl. (Orchidaceae) from Arunachal Pradesh, India

A. Nageswara Rao

Orchid Research Centre, State Forest Research Institute, Tipi, Bhalukpong - 790 114, Arunachal Pradesh, India

Abstract

The concept of treating the orchid genus *lone* Lindl. separate from *Sunipia* Lindl. is supported. A new species of Orchidaceae viz., *lone arunachalense* Nageswara Rao is described and an identification key is provided for the 6 species of *lone* found in Arunachal Pradesh.

INTRODUCTION

Lindley established two genera in the family Orchidaceae viz. Sunipia and lone in 1833 and 1853 respectively based on the structure of pollinarium. According to him, the former has pollinia with two free stipes with 2 seperate viscidia while the latter has pollinia with 2 stipes which are fused at apex and attached to a single common viscidium.

Later Hooker (1890) accepted Sunipia Lindl. as a separate genus but denied the genus status to Ione Lindl. and transferred all the species of Ione to the genus Bulbophyllum Thou. King and Pantling (1898) while describing the orchids of Sikkim recognised the species of Sunipia and Ione as quite distinct from Bulbophyllum, but however considered the former two as congeneric and surprisingly preferred to retain the name Ione as a valid name over earlier published name Sunipia and considered the latter as only a "legendary genus." This has been followed by Seidenfaden (1969) in his monographic work on the genus Ione Lindl.

Subsequently, Hunt (1971) recognised the priority of *Sunipia* over *Ione* and transferred all the species of *Ione* to *Sunipia*, treating *Ione* as a synonym. This was followed by later orchid taxonomists like Pradhan (1979), Seidenfaden (1986) et al.

Now, of late, orchidologists like Garay et al. (1994) and Seidenfaden (1995) have realized Lindley's old concept and recognised *lone* and *Sunipia* as two distinct genera.

During the course of critical study on the species of *lone Lindl*. in Arunachal Pradesh, a specimen from Tale Valley in Subansiri district was found to be quite distinct from the so far known species of the genus and hence it is described here as a new ones.

A. Nageswara Rao

Ione arunachalense Nageswara Rao, sp. nov. (Fig. 1)

lone minor Seidenf, affinis, sed differt foliis petiolarus flore singulari, sepalis lateralia liberis, labello lobo intermedio acuminato.

Type: India, Arunachal Pradesh, Lower Subansiri district, Tale Valley (1200 m), *Ingalhalli & Bennedict* 20350 (Holo-Orchid Herbarium, Tipi).

Small epiphytic herb. Rhizome ca. 1 mm thick, slender. Pseudobulbs 6-9 x 5-8 mm, ovoid, ca. 2 cm apart on the rhizome with slender roots below, 1-leaved. Leaves ca. 3 x 0.5 cm, oblong, subacute, with a distinct petiole of ca. 7 mm long. Scape ca. 2 cm long, arising from the base of the pseudobulb. Flower solitary, ca. 2 cm long, crimson coloured. Bract ca. 6 x 1 mm, lanceolate, acute, shorter than the pedicellate ovary. Sepals ca. 10 x 2.5 mm, subequal, lanceolate, acute, 3-nerved; lateral sepals free. Petals ca. 3 x 3 mm, ovate-suborbicular, minutely apiculate, 1-nerved. Lip about 9 mm long, with a median narrow ridge on the disc from base to the middle; lateral lobes 3 x 5 mm, auricular, entire; midlobe ca. 6 mm long, linear, acuminate. Column very short, ca. 1 mm long. Anther small, 2-celled. Pollinia 4 in 2 groups, each group with a short, linear stipe and a gland at apex. Rostellum with 2 processes in front. Ovary with pedicel ca. 12 mm long, pale green.

Flowering: June

Distribution: Known only from lower Subansiri district, Tale Valley (1200 m), in wet evergreen subtropical forest.

Ione arunachalense Nageswara Rao sp. nov, is very close to I. minor Seidenf. which is an endemic species of Thailand but differs from it in having petioled leaves, single-flowered inflorescence, free lateral sepals and linear-acuminate midlobe of lip.

With the present new species the genus *lone* is represented in Arunachal Pradesh by six species which may be identified by the following key:

- 1. Lateral sepals free.
 - 2. Flowers solitary.

 - 3. Scape more than 1 cm long; flowers small; sepals 1 cm long I. arunachalense

Ione arunachalense sp. nov. (Orchidaceae)




Fig. 1. *Ione arunachalense* Nageswara Rao, *sp. nov.* A. Habit; B. Dorsal sepal; C. Lateral sepal; D. Petal; E. Lip; F. Column with pollinia *in situ* (Drawn from the holotype-*Ingalhalli et Bennedict* 20350).

A. Nageswara Rao

- - 4. Petals with serrate margins; lip broader at basal half and narrowed at apex.

1. Lateral sepals connate or coherent along the inner margins.

- 5. Inflorescene upto 10 cm long; flowers erect, bracts not scarious, petals oblong

 I. candida

Acknowledgements

Sincere thanks are due to Mr. G.P. Shukla, IFS, Principal Chief Conservator of Forests; Mr. K. Namchoom, IFS, Chief Conservator of Forests and Mr. M.L. Deori, IFS, Director, State Forest Research Institute, Arunachal Pradesh for encouragement and facilities. Thanks are also due to Dr. Gunnar Seidenfaden, Denmark for his valuable comments on the new taxon.

Literature cited

Garay, L. S., F. Hamer & E. S. Siegerist. 1994. The genus *Cirrhopetalum* and the genera of the *Bulbophyllum* alliance. *Nord. J. Bot.* 14: 609-646.

Hooker, J. D. 1890. The Flora of British India. 5: 667-858.

Hunt, P. F. 1971. Notes On Asiatic orchids - 6. Kew Bull. 26: 171-185.

King, G. & R. Pantling. 1898. The orchids of Sikkim Himalaya. Ann. Roy. Bot. Gard. Calcutta 8: 1-342.

Lindley, J. 1830-40. The genera and species of Orchidaceae Plants. London.

Lindley, J. 1852-59. Folia Orchidaceae. London.

Pradhan, U. C. 1979. Indian orchids - Guide to identification and culture. Vol. 2. Faridabad.

Seidenfaden, G. 1969. Notes on the genus lone. Bot. Tidsskr. 64(2-3): 205-238.

Seidenfaden, G. 1986. Orchid Genera in Thailand xiii - Thirty three epidendroid genera. *Opera Bot.* 89: 1-214.

Seidenfaden, G. 1995. Contribution to the orchid flora of Thailand XII. Opera Bot. 124: 1-90.