


SEM-characterisation of achene morphology towards the taxonomy of Indian species of *Saussurea* DC. (Asteraceae)

Saklani A., Rao R.R. & L.B. Chaudhary



How to cite:

Saklani A., Rao R.R. & L.B. Chaudhary 2000. SEM-characterisation of achene morphology towards the taxonomy of Indian species of *Saussurea* DC. (Asteraceae). *Rheedea* 10(1): 1–18.

<https://dx.doi.org/10.22244/rheedea.2000.10.01.01>

Published in print: 30.06.2000

Published Online: 01.01.2022

SEM-characterisation of achene morphology towards the taxonomy of Indian species of *Saussurea* DC. (Asteraceae)

A. Saklani, R. R. Rao and L. B. Chaudhary

Taxonomy & Biodiversity Division, National Botanical Research Institute
Lucknow - 226 001, India

Abstract

The paper deals with SEM-characterisation of achenes of 23 species of *Saussurea* DC. in India. The micromorphological characters of achenes are highly useful in the identification of different species of the genus. A key for identification of 23 species of *Saussurea* based on achene characters is provided.

INTRODUCTION

The genus *Saussurea* DC. comprises about 300 species in the world (Bremer, 1994) and about 61 species in India (Hajra, 1988, 1995). The genus is chiefly distributed in temperate Asia, with one species each in Europe, Australia and North and West America. *Saussurea* DC. is characterised by spineless leaf margins; usually purple or bluish flower heads with only disk florets; spineless multiseriate involucral bracts; receptacle with bristly scales; usually bi-seriate pappus, the outer usually of rigid bristles and the inner longer plumose with their bases fused in a ring.

The identification of some species of the genus is problematic due to their morphological plasticity. The distinction between the various *Saussurea* species is normally based on macromorphological parameters like habit and size of plants, size and shape of leaves and capitula, and the nature of phyllaries (Clarke, 1876; Hooker, 1881; Lipschitz, 1979; Hajra, 1988).

Many workers (Martin & Barkley, 1961; Dittrich, 1966, 1968, 1970; Kynclova-Petrocyova, 1970; Borgen, 1972; Singh *et al.*, 1972; Rajashekhar & Razi, 1976-77; Wagenitz, 1976; Chaudhary & Pandey, 1995) have considered the achene characters for segregation of species. According to Kynclova-Petrocyova (1970) the general shape of the achenes, the presence or absence of wings, carinae, ribs and border, and the colour are all reliable and stable distinguishing characters. Dittrich (1968) while examining the morphological characters of achenes of subtribe *Cardueae* - *Centaureinae* found the shape of the hilum, detachment of the

A. Saklani *et al.*




Fig. 1. Scanning electron micrographs of achenes of *Saussurea*

Achene morphology of *Saussurea* DC.

fruits, presence of the epicarp and also the hairiness of the fruit as important characters for the identification of genera. Dittrich (1970) also investigated the fruits of 17 genera including *Saussurea* belonging to subtribe *Cardueae-Carduinae* and gave a key to the genera based on the fruit characters. Dittrich (1970) accepted that *Saussurea* stands apart in several important features and is taxonomically isolated from the other genera.

The present study deals with the morphology of the achenes of 23 species of *Saussurea* under the light as well as Scanning Electron Microscope with a view to explore more characters of individual taxa and to assess and establish the taxonomic significance of them. The characters such as the base, the surface and the apex of achenes were found useful in differentiating the species and as such a key to all species studied established. The pappus bristles do not exhibit much variations among different species except *S. yakla* (Fig. 7, A), *S. simpsoniana* (Fig. 7, B), *S. albescens* (Fig. 7, C) and therefore, have not been discussed.

MATERIALS AND METHODS

Achenes of 23 species of *Saussurea* gathered afresh from the field as well as from the herbarium specimens have been examined under the light microscope and Scanning Electron Microscope (SEM). For SEM studies, the dry cleaned matured achenes were gold coated with SCD sputter coater and finally observations were carried out with a Phillips SEM XL20 at 15-20 KV and photographs were taken. For each species at least 4-5 achenes from different populations were scanned but photographs were taken only of the specimens cited under each species. Herbarium acronyms used are those given by Holmgren *et al.* (1990).

OBSERVATIONS

Saussurea abnormis Lipsch. (Fig. 1, A&B)

Achenes 5.0 x 1.2 mm, narrowly obovate, contracted at apex, apex truncate, hilum basal, surface reticulate.

Specimen examined: Uttar Pradesh, Chamoli Garhwal, opposite Niti village, 3800 m, Naithani 56137 (BSD).

Saussurea albescens (DC.) Sch.-Bip. (Fig. 1, C&D)

Achenes 6.4 x 2.0 mm, narrowly obovate, straight, apex truncate and wavy, some irregular spiny structures below the apex (neck portion), hilum basal, surface rugose/muricate.

Explanation to Fig.1. A & B. *S. abnormis*: A. Achene, B. Surface of achene; C & D. *S. albescens*: C. Achene, D. Apical portion of the achene; E & F. *S. auriculata*: E. Achene, F. Surface of achene; G. *S. bracteata*: surface of achene; H. *S. candelleana*: Achene (Scale bars: A, C, E & H = 1 mm; B & G = 50 µm; F = 20 µm).

A. Saklani *et al.*




Fig. 2. Scanning electron micrographs of achenes of *Saussurea*

A. Saklani *et al.*




Fig. 3. Scanning electron micrographs of achenes of *Saussurea*

Achene morphology of *Saussurea* DC.

Specimen examined: Himachal Pradesh, Bashahr, Sarahan chaur road, Raizada 19800 (DD).

***Saussurea auriculata* (Spreng. ex DC.) Sch.-Bip. (Fig. 1, E&F)**

Achenes 3.6 x 1.0 mm, 4-angled, narrowly obovate, straight, laciniate at apex, hilum lateral, surface sparsely spiny at lower magnification, ridged at higher magnification.

Specimen examined: Uttar Pradesh, Uttar Kashi, on way to Harkidun, 3000 m, Rana & Saklani 210142 (LWG).

***Saussurea bracteata* Decne. (Fig. 1, G)**

Achenes 4.5 x 1.0 mm, narrowly obovate, slightly curved, apex truncate, hilum sub-lateral, surface striate.

Specimen examined: Uttar Pradesh, Kumaon, Dhauli valley, Duthie 3038 (DD).

***Saussurea candolleana* Wall. ex DC. (Figs. 1, H; 2, A & B)**

Achenes 2.5 x 1.0 mm, 4-5-angled, oblong, straight, slightly contracted in middle, apex truncate, hilum basal, surface faintly rugose.

Specimen examined: Uttar Pradesh, Kumaon, Pindari, 3500-5000 m, Rao 4581 (BSD).

***Saussurea costus* (Falc.) Lipsch. (Fig. 2, C)**

Achenes 8.5-11.0 x 2.5-4.0 mm, obovate, straight, apex truncate, hilum basal, surface striate.

Specimen examined: Uttar Pradesh, Uttarkashi, Barsu, 2500 m, Saklani 216699 (LWG).

***Saussurea fastuosa* (Decne.) Sch.-Bip. (Fig. 2, D&F)**

Achenes 6.4 x 1.3 mm, oblong, slightly curved, apex truncate with a club-shaped beak, hilum basal, surface striate.

Specimen examined: Uttar Pradesh, Chamoli Garhwal, Valley of flowers, 3500 m, Rao 31743 (BSD).

Explanation to Fig. 2. A & B. *S. candolleana*: A. Apical portion of achene, B. Surface of achene; C. *S. costus*: Surface of achene; D, E & F. *S. fastuosa*: D. Achene, E. Apical portion of the achene, F. Surface of achene; G & H. *S. glacialis*: G. Achene, H. Surface of achene (Scale bars: A & E = 200 µm; B, C, F & H = 50 µm; D = 2 mm; G = 1 mm).

Achene morphology of *Saussurea* DC.***Saussurea glacialis* Herd. (Fig. 2, G&H)**

Achenes 3.0 x 1.1 mm, oblong, slightly curved, apex like a pitcher head, hilum lateral, surface reticulate.

Specimen examined: Jammu & Kashmir, Ladakh, Thang Lang La, 5200 m, Bhattacharyya 41036 (BSD).

***Saussurea glanduligera* Sch.-Bip. ex Hook. f. (Fig. 3, A&B)**

Achenes 5.1 x 2.0 mm, 4-angled, obovate, slightly curved towards base, apex slightly concave, collar present, hilum sub-lateral, surface reticulate.

Specimen examined: Himachal Pradesh, Spiti, Gyundi Nala, 3900 m, Bhattacharyya 49377 (BSD).

***Saussurea gnaphalodes* (Royle ex DC.) Sch.-Bip. (Fig. 3, C&D)**

Achenes 3.8 x 1.2 mm, 5-angled, narrowly obovate, straight, apex obtuse, hilum lateral, surface ribbed.

Specimen examined: Jammu & Kashmir, Ladakh, Thang Lang La, 5000 m, Bhattacharyya 41031 (BSD).

***Saussurea graminifolia* Wall. ex DC. (Fig. 3, E&F)**

Achenes 3.3 x 1.3 mm, obovate, straight, apex obtuse, irregularly wavy ring at the base of apex, hilum basal, surface rugose.

Specimen examined: Uttar Pradesh, Pindari, Furkia, 3500-5000 m, Rao 4512 (BSD).

***Saussurea heteromalla* (D. Don) Hand.-Mazz. (Fig. 3, G&H; 4, A)**

Achenes 3.3 x 1.1 mm, 4-5-angled, obovate, straight, apex truncate and spiny, hilum lateral, surface muricate.

Specimen examined: Uttar Pradesh, Uttarkashi, Sankari, 1800 m, Rana & Party, 219718 (LWG).

Explanation to Fig. 3. A. & B. *S. glanduligera*: A. Achene, B. Surface of achene; C & D. *S. gnaphalodes*: C. Achene, D. Surface of achene; E & F. *S. graminifolia*: E. Achene, F. Surface of achene; G & H. *S. heteromalla*: G. Achene, H. Surface of achene. (Scale bars: A, C, E & G = 1 mm; B, D & F = 50 μ m; H = 500 μ m).

A. Saklani *et al.*




Fig. 4. Scanning electron micrographs of achenes of *Saussurea*

Achene morphology of *Saussurea* DC.

***Saussurea hookeri* C. B. Clarke (Fig. 4, B&C)**

Achenes 3.5 x 1.0 mm, almost oblong, slightly curved and narrow towards the base, apex truncate, hilum basal, surface striate.

Specimen examined: Sikkim, Lachung, G. A. Gammie s. n. (DD).

***Saussurea jacea* (Klotzsch) C. B. Clarke (Fig. 4, D)**

Achenes 8.0 x 1.5 mm, oblong, straight, apex truncate, hilum basal, surface ridged under light microscope.

Specimen examined: Jammu & Kashmir, Ladakh, Dras, 2700 m, Bhattacharyya 41198 (BSD).

***Saussurea leontodontoides* (DC.) Sch.-Bip. (Fig. 4, E,F&G)**

Achenes 5.0 x 1.3 mm, more or less oblong, straight, apex obtuse with some irregular structure, hilum basal, surface muricate in lower half at low magnification, striate at higher magnification.

Specimen examined: Uttar Pradesh, Chamoli Garhwal, Amritganga valley, 3000 m, Naithani 37981 (BSD).

***Saussurea obvallata* (DC.) Edgew. (Fig. 4, H & 5, A)**

Achenes 5.2 x 1.0 mm, oblong, straight, sometimes contracted in middle, apex truncate, hilum basal, surface reticulate.

Specimen examined: Uttar Pradesh, Chamoli Garhwal, Hemkund, 4200 m, Bhattacharyya 24489 (BSD).

***Saussurea piptathera* Edgew. (Fig. 5, B)**

Achenes 5.0 x 1.3 mm, oblong, straight, apex sub-truncate, hilum lateral, surface ridged.

Specimen examined: Himachal Pradesh, Lahul, Koksar, 3425 m, Bhattacharyya 17029 (BSD).

Explanation to Fig. 4. A. *S. heteromalla*: Surface of achene; B & C. *S. hookeri*: B. Achene, C. Surface of achene; D. *S. jacea*: Apical portion of achene; E, F & G. *S. leontodontoides*: E. Achene, F. Apical portion of achene, G. Surface of achene; H. *S. obvallata*: Achene (Scale bars: A, C & G = 50 µm; B, E & H = 1 mm; D & F = 500 µm).

A. Saklani *et al.*




Fig. 5. Scanning electron micrographs of achenes of *Saussurea*

Achene morphology of *Saussurea* DC.

***Saussurea roylei* (DC.) Sch.- Bip. (Fig. 5, C,D&E)**

Achenes 5.0 x 1.3 mm, narrowly obovate, straight, apex laciniate, hilum sub-lateral, surface faintly reticulate.

Specimen examined: Uttar Pradesh, Uttarkashi, Bhojwasa, 3700 m, Bhattacharyya 74761 (BSD).

***Saussurea simpsoniana* (Field. & Gard.) Lipsch. (Fig. 5,F)**

Achenes 2.5 x 1.1 mm, 4-5-angled, obovate, straight, apex rounded, hilum sub-lateral, surface rugose.

Specimen examined: Uttar Pradesh, Chamoli Garhwal, Hemkund, 4200 m, Rao 31771 (BSD).

***Saussurea stracheyana* (Kuntze) Lipsch. (Fig. 5, G&H)**

Achenes 3.3 x 0.7 mm, 4-5-angled, oblong, slightly curved, apex truncate, contracted at neck, hilum lateral, surface faintly reticulate.

Specimen examined: Uttar Pradesh, Uttarkashi, on way to Harkidun, 2460 m, Rana & Saklani 210137 (LWG).

***Saussurea taraxicifolia* Wall. ex Sch.-Bip. (Fig. 6, A&B)**

Achenes 4.5 x 0.8 mm, obovate, straight, apex truncate with papillose structure, hilum basal, surface rugose.

Specimen examined: Uttar Pradesh, Chamoli Garhwal, Keal Bhairawn, 3600 m, Kapoor & Party 70579 (LWG).

***Saussurea uniflora* (DC.) Wall. ex Sch.-Bip. (Fig. 6, C&D)**

Achenes 4.2 x 1.0 mm, oblong, slightly curved, apex obtuse, hilum basal and surface ribbed.

Specimen examined: Sikkim, Lachung, Gammie s. n., (DD).

Explanation to Fig. 5. A. *S. obvallata*: Surface of achene; B. *S. poptathera*: Surface of achene; C, D & E. *S. roylei*: C. Achene, D. Apical portion of the achene, E. Surface of achene; F. *S. simpsoniana*: Achene; G & H. *S. stracheyana*: G. Achene, H. Surface of achene (Scale bars: A & H = 20 µm; B & E = 50 µm; C, F & G = 1 mm; D = 500 µm).

A. Saklani *et al.*




Fig. 6. Scanning electron micrographs of achenes of *Saussurea*. A & B. *S. taraxifolia*: A. Apical portion of the achene; B. Basal portion of the achene; C & D. *S. uniflora*: C. Achene, D. Surface of the achene; E & F. *S. yakla*: E. Achene, F. Surface of achene (Scale bars: A = 500 µm; B = 200 µm; C & E = 1 mm; D & F = 50 µm).

Achene morphology of *Saussurea* DC.

Saussurea yakla C. B. Clarke (Fig. 6, E&F)

Achenes 7.7 x 2.0 mm, obovate (spatulate), almost straight, apex concave, hilum basal, surface beaded, cells faint, rectangular and arranged in linear rows.

Specimen examined: South East Lhasa, Chapman 113 (DD).

DISCUSSION AND CONCLUSIONS

The morphological characters such as the shape of the achene, the areole, the apex and surface of achenes are usually stable and reliable and hence they are taken into consideration for taxonomic treatment. Based on achene characters a key for the identification of 23 species of *Saussurea* is provided.

The size of achenes is not a reliable distinguishing character in most of the cases, however, in some species like *A. graminifolia*, *S. yakla*, *S. abnormis*, *S. costus* and *S. jacea*, it can be used as a key character. The length and width ratio of achenes is usually 3:1 to 4:1, but in *S. simpsoniana* and *S. jacea* it ranges from 2:1 to 5:1 respectively.

In most species, the achenes are obovate, narrowly obovate or oblong. Commonly the achenes are straight or sometimes slightly curved (*S. glacialis* and *S. hookeri*).

Among the species investigated, the hilum is basal in as many as 13 species, while in *S. glanduligera*, *S. bracteata*, *S. roylei* and *S. simpsoniana* it is sub-lateral, and in the species like *S. auriculata*, *S. glacialis*, *S. gnaphalodes*, *S. heteromalla*, *S. piptathera* and *S. stracheyana* the hilum is lateral.

The apex of achenes is usually truncate or obtuse, or sometimes concave as in *S. glanduligera* and *S. yakla*. In a few species such as *S. albescens*, *S. auriculata*, *S. heteromalla*, *S. neglecta* and *S. roylei* the apex is like a cup and the pappus arises inside the cavity. In certain species the margins of cup become lacinate (*S. roylei*), wavy (*S. albescens*), spiny (*S. heteromalla*) or divaricate (*S. leontodontoides*). In *S. taraxicifolia* the apex has some papillose structure. In *S. glacialis* pitcher's head like structure is formed at the apex while in *S. fastuosa* the apex is protruded into a beak-like structure.

At lower magnification the surface of achenes in most of the species is rugose, striate or smooth. In *S. heteromalla* the surface is muricated while it is tubercled in *S. auriculata*. The lower half of the achenes in case of *S. leontodontoides* has ornate patterns while the upper portion is almost striate. At higher magnification (above 500 x) under SEM many patterns on the surface such as reticulate, rugose, striate and ribbed are commonly observed. Sometimes the surface is ridged (*S. auriculata* and *S. piptathera*), muricate (*S. heteromalla*) or beaded (*S. yakla*).

A. Saklani *et al.*




Fig. 7. Scanning electron micrographs of pappus bristles in *Saussurea*. A. *S. yakla*; B. *S. simpsoniana*; C. *S. albescens*; D. *S. obvallata* (Scale bars: A, B & D = 200 μm ; C = 100 μm).

Achene morphology of *Saussurea* DC.

In most of the species of *Saussurea* studied, the pappus are plumose (feathery). The secondary pappus hairs on the inner pappus bristles are long and distributed throughout the surface. The surface of bristles is striate in all species investigated. However, at least 4 types of inner pappus bristles are found in the species investigated - (a) bristles with minute and sparsely distributed secondary pappus hairs throughout the surface (*S. yakla*, Fig. 7, A) (b) bristles with only a few long, secondary pappus hairs, that too sparsely distributed on their margins only (*S. simpsoniana*, Fig. 7, B), (c) bristles with longer and a few minute secondary pappus hairs sparsely distributed on the bristle surface (*S. albescens*, Fig. 7, C), and (d) the bristles with long feathery secondary pappus hairs, common in rest of the species (Fig. 7, D).

Key to the species based on achene morphology

- | | | |
|-----|--|-------------------------|
| 1a. | Surface of achenes spiny | 2 |
| 1b. | Surface of achenes not spiny | 3 |
| 2a. | Achenes laciniate at apex | <i>S. auriculata</i> |
| 2b. | Achenes wavy (spiny) at apex | <i>S. heteromalla</i> |
| 3a. | Hilum of achenes basal | 4 |
| 3b. | Hilum of achenes lateral or sub-lateral | 16 |
| 4a. | Achenes narrowly obovate to obovate | 5 |
| 4b. | Achenes oblong | 10 |
| 5a. | Apex of achenes obtuse to concave | 6 |
| 5b. | Apex of achenes truncate | 7 |
| 6a. | Achenes 3.3 x 1.3 mm, obtuse with irregular wavy ring at apex, surface rugose | <i>S. graminifolia</i> |
| 6b. | Achenes 7.7 x 2.0 mm, concave at apex, surface beaded | <i>S. yakla</i> |
| 7a. | Surface of achenes rugose | 8 |
| 7b. | Surface of achenes reticulate or smooth | 9 |
| 8a. | Apex of achenes wavy with some irregular spiny structure around the neck | <i>S. albescens</i> |
| 8b. | Apex of achenes papillose | <i>S. taraxicifolia</i> |
| 9a. | Achenes 5.0 x 1.2 mm, narrowly obovate, contracted at apex, surface reticulate | <i>S. abnormis</i> |
| 9b. | Achenes 8.5-11.0 x 2.5-4.0 mm, obovate, surface smooth and black spotted | <i>S. costus</i> |

A. Saklani et al.

- 10a. Achenes with club-shaped beak *S. fastuosa*
- 10b. Achenes with obtuse or truncate apex 11
- 11a. Apex of achenes obtuse 12
- 11b. Apex of achenes truncate 13
- 12a. Achenes oblong, straight, surface muricate to striate in lower half ... *S. leontodontoides*
- 12b. Achenes oblong, slightly curved, surface ribbed *S. uniflora*
- 13a. Surface of achenes ridged *S. jacea*
- 13b. Surface of achenes almost smooth 14
- 14a. Achenes reticulate *S. obvallata*
- 14b. Achenes striate or faintly rugose 15
- 15a. Achenes 4-5-angled, surface faintly rugose *S. candolleana*
- 15b. Achenes almost terete, slightly curved, surface striate *S. hookeri*
- 16a. Apex of achenes pitcher head like *S. glacialis*
- 16b. Apex of achenes otherwise 17
- 17a. Achenes laciniate at apex *S. roylei*
- 17b. Achenes obtuse, rounded, subtruncate or concave at apex 18
- 18a. Apex of achenes obtuse or rounded 19
- 18b. Apex of achenes sub-truncate, truncate, or slightly concave 20
- 19a. Achenes with obtuse, lateral hilum, surface ribbed *S. gnaphalodes*
- 19b. Achenes with round apex, sub-lateral hilum, rugose surface *S. simpsoniana*
- 20a. Apex of achenes slightly concave, with a distinct collar *S. glanduligera*
- 20b. Apex of achenes truncate or sub-truncate, collar absent 21
- 21a. Achenes 4-angled, contracted at neck, surface faintly reticulate *S. stracheyana*
- 21b. Achenes almost terete, surface striate or ridged 22
- 22a. Surface of achenes striate, hilum sub-lateral *S. bracteata*
- 22b. Surface of achenes ridged, hilum lateral *S. piptathera*

Achene morphology of *Saussurea* DC.

Acknowledgements

The authors express their thanks to the Director, National Botanical Research Institute, Lucknow for the facilities provided. Thanks are also due to the Directors/Scientists-in-Charge of the herbaria BSI and DD for giving permission to examine the herbarium specimens. One of the authors (A. S.) is thankful to Council of Scientific and Industrial Research, New Delhi for awarding him Research Associateship.

Literature Cited

- Borgen, L. 1972. Embryology and achene morphology in endemic Canarian species of *Chrysanthemum* (L.) Hoffm. sub-genus *Argyranthemum* (Webb.) Harling (Asteraceae). *Norw. J. Bot.* 19 (3-4): 149-170.
- Bremer, K. 1994. *Asteraceae - cladistic and classification*. Timber Press, Portland, Oregon, U. S. A..
- Chaudhary, L. B. & A. K. Pandey. 1995. Taxonomic significance of phyllaries and achenes of *Tricholepis* DC. (Asteraceae). In: A. K. Pandey (Ed.), *Taxonomy and Biodiversity*. CBS Publishers and Distributors, New Delhi. pp. 164-171.
- Clarke, C.B. 1876. *Compositae Indicae descriptae et secus genera Benthamii Ordinatae*. Calcutta.
- Dittrich, M. 1966. Karpologische untersuchungen zur syetmatik von *Centaurea* und verwandten Gattungen. *Bot. Jb. Syst.* 88: 70-162.
- Dittrich, M. 1968. Morphologische untersuchungen an fruchten der subtribus Cardueae-Centaureinae (Compositae). *Willdenowia* 5: 67-107.
- Dittrich, M. 1970. Morphologische und anatonomische untersuchungen an fruchten der Carduinae (Compositae). I. morphologischer teil. *Candolleana* 25 (1): 45-67.
- Hajra, P. K. 1988. *Brahmkamal and its Allies (Saussurea the genus)*. Jugal Kishore & Co., Dehra Dun, India.
- Hajra, P. K. 1995. Tribe Cardueae. In: P. K. Hajra , R. R. Rao, D. K. Singh & B. P. Uniyal (Eds.), *Flora of India*. Botanical Survey of India, Calcutta. Vol. 12: 150-226.
- Holmgren, P. K., N. H. Holmgren & L. C. Barnett. 1990. *Index Herbariorum. Part I: The Herbaria of the World* (8th Ed.), Utrecht.
- Hooker, J. D. 1881. Compositae. In: J. D. Hooker, *Flora of British India*. L. Reeve & Co., London. Vol.3: 219-419.
- Kynclova-Petrocyora, M. 1970. Comparative morphology of achenes of the tribe Anthemidae Cass. (Family Asteraceae) and its taxonomic significance. *Perslia (Praha)* 42: 33-53.
- Lipschitz, S. J. 1979. *Rod Saussurea DC. (Asteraceae)*. Leningrad.

A. Saklani *et al.*

- Martin, A. C. & W. D. Barkley. 1961. *Seed Identification Manual*. Univ. California Press, Berkeley & Los Angeles.
- Rajashekara, G. & B.A. Razi. 1976-77. Studies on fruits of Asteraceae of Mysore city as an aid to classification. *J. Mysore Univ. Sect. -B. Sci.* 27: 197-211.
- Singh, G., B. Misri & P. Kachroo 1972. Achene morphology: an aid to taxonomy of Indian plants. 1. Compositae, Liguliferae. *J. Ind. Bot. Soc.* 51: 235-242.
- Wagenitz, G. 1976. Systematics and Phylogeny of the Compositae (Asteraceae). *Pl. Syst. Evol.* 125: 29-46.