


# Cleome burmannii (Cleomaceae): A New Record to the Flora of Rajasthan, India

Yadav B.L. & K.L. Meena



How to cite:

Yadav B.L. & K.L. Meena 2009. *Cleome burmannii* (Cleomaceae): A New Record to the Flora of Rajasthan, India. *Rheedea* 19(1&2): 65-66.

https://dx.doi.org/10.22244/rheedea.2009.19.01.16

Received: 09.06.2008 Revised and accepted: 20.07.2009

Published in print: 30.12.2009 Published Online: 30.12.2009



This volume of Rheedea is published with partial financial assistance from Department of Science and Technology, Government of India, New Delhi



# Cleome burmannii (Cleomaceae): A New Record to the Flora of Rajasthan, India

B. L. Yadav\* and K. L. Meena

Department of Botany, MLV Government College, Bhilwara – 311 001, Rajasthan, India.

\*E-mail: drblyadav@yahoo.com

#### **Abstract**

The taxon *Cleome burmannii* Wight & Arn. (Cleomaceae) so far known from South India, Sri Lanka and Indonesia is reported for the first time from Rajasthan, India.

Keywords: Cleome burmannii, New Record, Rajasthan, India

# Introduction

Genus *Cleome* L. distributed in the tropical America and Africa comprises *c.* 150 species (Mabberley, 2005) with 15 wild species in Indian flora (Sundara Raghavan, 1993). Bhandari (1978), Sharma & Tiagi (1979) and Shetty & Singh (1987) have reported 4, 3 and 8 species of this genus respectively from Rajasthan.

We collected a few interesting specimens belonging to the family Cleomaceae in Bhilwara district of Rajasthan. The specimens were found allied to *C. rutidosperma* DC. but differed in certain floral characters. Critical study of these specimens and perusal of relevant literature (Babu & Majumdar, 1976; Kumar & Bir Bahadur, 1978; Sundara Raghavan, 1984) revealed that the identity of the specimens as *C. burmannii* Wight & Arn., a new record for Rajasthan.

Cleome burmannii Wight & Arn., Prodr. Fl. Ind. Orient. 1: 22. 1834; Hook.f. & Thomson in Hook.f., Fl. Brit. India 1: 170. 1872; Sundraragh. in B. D. Sharma & N. P. Balakr., Fl. India 2: 304, f. 57. 1993.

Flowering & Fruiting: May – November.

*Distribution*: India, Indonesia (Java) and Sri Lanka. India: Kerala, Tamil Nadu; now in Rajasthan.

Specimens Examined: INDIA, Rajasthan, Bhilwara District, Pansal, Meena & Yadav 51; Kotary, Meena & Yadav 221 (Herbarium, MLV Govt. College, Bhilwara).

Ecological Notes: Extremely rare in waste lands, sites of old cultivation and road sides on moist sandy soil and wet places up to 600 m altitude. The taxon grows in association with Cleome viscosa L., Dactyloctenium aegyptium (L.) Willd., Indigofera linnaei

Ali, *Polygala erioptera* DC., *Sida cordata* (Burm.f.) Borss.Waalk. and *Trianthema portulacastrum* L. The taxon has not been recorded in similar habitats elsewhere in Bhilwara district of Rajasthan inspite of repeated searches for the past five years. It is restricted to *c.* 2 sq. km. area and is represented by approximately 100 individuals. The occurrence of this species in Gujarat and some reports on the occurrence from Kerala and Tamil Nadu are based on misidentifications (Sundara Raghavan, 1984, 1993). The disjunct distribution of the species in India is of phytogeographical interest. This species has been evaluated as Endangered in India.

## Acknowledgements

Authors are grateful to Dr. M. Sanjappa, Director and Dr. M. S. Mondal, Joint Director, Botanical Survey of India, Howrah, for confirming the identity of the specimens. Thanks are also due to Dr. U. C. Bansal, Principal, Dr. D. C. Jain, Dr. V. K. Dosi and Dr. S. S. Bhatt, Vice-Principals, MLV Government College, Bhilwara, for providing the facilities. Financial assistance provided by UGC, Bhopal, is gratefully acknowledged.

### Literature Cited

Babu, C. R. & N. C. Majumdar 1976. Taxonomical Notes on *Cleome aspera* Koen. ex DC., *C. burmannii* Wight & Arn. and *C. rutidosperma* DC. (Cleomaceae). *J. Bombay Nat. Hist. Soc.* 71: 629 – 632.

**Bhandari, M. M. 1978.** *Flora of Indian Desert*. MPS. Repros, Jodhpur.

- Kumar, P. V. & Bir Bahadur 1978. Seed morphology of thirteen species of Cleome L. (Capparidaceae). J. Indian Bot. Soc. 57: 39 - 46.
- Mabberley, D. J. 2005. The Plant Book: A Portable Dictionary of the Vascular Plants. (South Asian Edition). Cambridge University Press, Cambridge, UK.
- Sharma, S. & B. Tiagi 1979. Flora of North-East Rajasthan. Kalyani Publishers, New Delhi.
- Shetty, B. V. & V. Singh 1987. Flora of Rajasthan. Vol. 1. Botanical Survey of India, Howrah.

- Sundara Raghavan, R. 1984. On Cleome burmannii Wt. & Arn. (Capparaceae) - Its identity and distribution. J. Econ. Taxon. Bot. 5: 463 – 465.
- Sundara Raghavan, R. 1993. Capparaceae. In: Sharma, B. D. & N. P. Balakrishnan (Eds), Flora of India. Vol. 2. Botanical Survey of India, Howrah. pp. 248 – 335.

Received: 9.6.2008

Revised and Accepted: 20.7.2009