


Rhynchosia hainesiana Satyan. et Thoth. (Fabaceae)- A new record for Southern Peninsular India

Sri Rama Murthy K., Pullaiah T. & S. Sandhya Rani



How to cite:

Sri Rama Murthy K., Pullaiah T. & S. Sandhya Rani 1999. *Rhynchosia hainesiana* Satyan. et Thoth. (Fabaceae)- A new record for Southern Peninsular India. *Rheedea* 9(1): 37–39.

https://dx.doi.org/10.22244/rheedea.1999.09.01.04

Published in print: 30.06.1999 Published Online: 01.01.2022





Rhynchosia hainesiana Satyan. et Thoth. (Fabaceae)-A new record for Southern Peninsular India

K. Sri Rama Murthy, T. Pullaiah and S. Sandhya Rani

Department of Botany, Sri Krishnadevaraya University, Anantapur - 515003, Andhra Pradesh, India

Abstract

Rhynchosia hainesiana Satyan. et Thoth. is reported for the first time from Southern Penninsular India. Detailed description, illustration and relevant notes of this taxon are given.

INTRODUCTION

The genus *Rhynchosia* Lour. with about 300 species (Santapau & Henry, 1973) is distributed in tropical and sub-tropical regions. Sanjappa (1991) reported 25 species of *Rhynchosia* from India. Recently van der Maesen (1995) added another species viz., *Rhynchosia courtallensis*, by transferring South Indian *Dunbaria latifolia* Wight & Arn.

While studying the Faboideae of Eastern Ghats, we collected an interesting specimen of *Rhynchosia* from Geddamannugu Kondur, Krishna district in Andhra Pradesh. After a critical study of specimen and relevant literature, it was identified as *Rhynchosia hainesiana* Satyan. Thoth. So far it is reported only from Angul forests of Orissa by Satyanarayana and Thothathri (1986). The taxon is described and illustrated here.

Rhynchosia hainesiana Satyan. et Thoth. in Bull. Bot. Surv. India 28: 243. 1986.

Erect undershrub, up to 75 cm height, branchlets erect, minutely hirsute with stiff, spreading, white canescent hairs, resinous gland-dotted. Stipules 1-1.5 mm long, linear, persistent, pubescent, exstipellate. Leaves 3-foliolate, 1-3 cm long, leaflets narrowly lanceolate, 0.6-1.6 x 0.1-0.5 cm, base cuneate, apex acute, margin recurved (folded towards lower surface), greenish with glandular hairs above, white silky with minute hairs beneath, petiole 4-8 mm long, petiolule 0.5-1mm. Racemes axillary, to 1 cm long, peduncle 3-5 mm long; Flowers ca 8 mm across, pedicel very slender, 2-3 mm long, grey pubescent. Bracts 1 mm long, subulate, densely canescent, persistent. Calyx campanulate, tube ca 1 mm long, lobes unequally divided, to 3 mm long, pubescent with glandular hairs. Corolla yellow, standard orbicular, ca 5 x 4 mm, rounded at apex, claw ca 1 mm, auricles very minute; wings ca 4 x 1 mm, narrowly obovate, rounded at apex; keels ca 4 x 1.5 mm, falcate, rounded at apex. Stamens diadelphous (9+1), staminal sheath ca 5 x 1.5 mm; filaments 1-1.5 mm long; anthers uniform, ca 0.4 x 0.2 mm. Ovary glabrous, 2-ovulate, ca 1.7 mm

K. Sri Rama Murthy et al.




Fig 1. Rhynchosia hainesiana Satyan. et Thoth.: A. Twig; B. Leaf; C. Calyx split open; D. Standard petal; E. Wing petals; F. Keel petals; G. Stamens; H. Pistil.

Rhynchosia hainesiana

long; style ca 6 mm long, curved, slightly broadened above, stigma capitate. Pods ca 1 x 0.5 cm, oblong, glabrescent, beak incurved, 2-seeded.

Fls. & Frs. : March-April.

Distribution: Andhra Pradesh and Orissa - endemic.

Specimens examined: Geddamannugu Kondur; Krishna district, KSM 17638.

Acknowledgements

The first author (KSM) is thankful to CSIR, New Delhi for the award of SRF.

Literature cited

Sanjappa, M. 1991. Legumes of India, Bishen Singh Mahendra Pal Singh, Dehra Dun.

Santapau, H. & A.N. Henry. 1973. A Dictionary of the Flowering plants in India, New Delhi.

Satyanarayana, P. & K. Thothathri. 1986. Three New species of *Rhynchosia* Lour. (Fabaceae) from India. Bull. Bot. Surv. India. 28: 241-246.

van der Maesen, L.J.G. 1995. *Rhynchosia courtallensis* van der Maesen a new name for the South Indian *Durbaria latifolia* Wt. & Arn. (Leguminosae-Papilionoideae), Rheedea, 5 (1): 54-59.



