




## Floristic diversity in the Neogene and Modern forests of Pondicherry - Neyveli area of Tamil Nadu - its impact on climatic and phytogeographic aspects

Ramanujam C.G.K.



How to cite:

**Ramanujam C.G.K. 1996.** Floristic diversity in the Neogene and Modern forests of Pondicherry - Neyveli area of Tamil Nadu - its impact on climatic and phytogeographic aspects. *Rheedeia* 6(1): 29–41.

<https://dx.doi.org/10.22244/rheedeia.1996.06.01.03>

*Published in print:* 30.06.1996

*Published Online:* 01.01.2022

## **Floristic diversity in the Neogene and Modern forests of Pondicherry–Neyveli area of Tamil Nadu – its impact on climatic and phytogeographic aspects**

**C. G. K. Ramanujam**

Palaeobotany-Palynology Laboratory,  
Department of Botany  
Post-Graduate College of Science,  
Saifabad, Hyderabad-500 004, India.

---

### **Abstract**

The Neogene plant fossil (mega and micro) preserved abundantly in the sandstones around Pondicherry and the lignite deposits near Neyveli in the Cauvery basin of Tamil Nadu have been extensively studied over the last three decades. On the whole, 108 genera referable to 53 families of Angiosperms could be recognised in the Neogene flora of this region. Arecaceae is the predominant monocot family. Anacardiaceae, Barringtoniaceae, Clusiaceae, Combretaceae, Dipterocarpaceae and Fabaceae are numerically better represented among the dicots. The totality of the Neogene floral spectrum reveals the occurrence of discrete pockets of estuarine swamps at the coast line and tropical moist evergreen forest away from the coastal belt.

As against the tropical wet evergreen Neogene vegetational type, the present day Pondicherry-Neyveli area shows only few strands of vegetation consisting of scrub-woodlands or the so-called dry evergreen forests. The probable reasons for this dramatic change in the climate and vegetational types in the Neogene *vis a vis* the present day, along with the phytogeographic significance of some taxa are highlighted.

### **INTRODUCTION**

The sandstone outcrops around Pondicherry and the vast lignite deposits at Neyveli are verily replete with a teeming multitude of varied plant fossils such as silicified and carbonized woods, mummified leaves and twigs and a galaxy of spore and pollen types of bewildering morphographic diversity. Over the last three decades these have been studied extensively resulting in a vast information that merits critical analysis and assessment of its relevance to our understanding of the Upper Tertiary vegetational complexes and climatic patterns.

Notwithstanding the great significance of the available knowledge of the Neogene Angiosperms *vis a vis* their modern counterparts, it is consistently glossed over by the students of extant plants. Among other things, this is attributable to the general apathy of the protago-

## C. G. K. Ramanujam

sists of Neobotany towards palaeobotanical data. The relationships of the Tertiary plant fossil with the modern taxa are inversely proportional to their geological age. The present survey utilizes collectively the data obtained from both palaeobotanical and palynological studies for a meaningful perspective of the Neogene floral assemblages.

## GEOLOGICAL SET UP OF PONDICHERRY — NEYVELI AREA

Table 1 provides information on the general geological succession of the area. The sedimentary regime spreading from Pondicherry to south of Rameswaram towards the Coromandel coast of Tamil Nadu represents the Cauvery main basin. The Pondicherry - Neyveli area lies in the Ariyalur-Pondicherry sub-basin. Detailed geological and geophysical investigations of this area were earlier carried out by Balasundar (1968), Kailasam (1968), Ramanathan (1968, 1979), Subramanyam (1969) and Gowrisankaran *et al.* (1987). The basement Precambrian horizon is succeeded by fossil-rich limestone, calcareous sandstone and marl of Ariyalur Formation (Upper Cretaceous). Highly fossiliferous marine Eocene sediments are found overlying the Ariyalur Formation in several borewells drilled in this sub-basin. The Mio-Pliocene Cuddalore Formation consisting of sandstones, clayey sandstones, sandy clay, and pockets of lignite, overlies unconformably the Eocene sediments. The Recent alluvium occupying the major part of the sub-basin tops the sequence (Table 1).

**Table 1. General geological succession in the Pondicherry-Neyveli area**  
(After N. G. K. Murthy, 1968; Subramanyan, 1969)

Age	Formation	Lithology
Recent and subrecent		Soils, coastal sands, alluvium, laterite, kankar
Miocene (or Mio-Pliocene)	Cuddalore Formation	Argillaceous sandstone, lignite, pebble-bearing sandstone, grits, sands, clays
----- Unconformity -----		
Eocene		Black clays, calcareous sandstones, fossiliferous limestone
----- Unconformity -----		
Late Cretaceous	Ariyalur Formation	Shell limestones, shales, marl
----- Unconformity -----		
Precambrian	Intrusives	Dolerites, pegmatites, charnockite, granitoid gneisses etc.

### Floristic diversity in the Neogene and Modern forests

The sandstones around Pondicherry are coarse-grained or almost gritty, mottled and argillaceous with a generous dosage of ferruginous concretions and contain a profusion of silicified woods, many of them of respectful dimensions. The lignite embedded in the sandstone and seen essentially as one major seam in the vicinity of Neyveli shows rich and varied plant debris at various horizons.

Except in the environs of Tiruvakkarai about 13 miles WNW of Pondicherry, with an abundance of coniferous woods, the rest of the localities in the area show almost exclusively a wealth of silicified angiosperm woods. The coniferous woods at Tiruvakkarai are podocarpaceous and constitute virtually a pure strand of *Podocarpus*. Whether these podocarpaceous remains were from local floristic scenario or got transported from nearby hilly tracts seems to be a moot point.

The source material for the lignite deposit at Neyveli, however, was made up exclusively of angiosperm taxa with only a sprinkling of pteridophytic elements.

The present study is based essentially on angiosperm floristics in view of the overwhelming numerical preponderance of angiosperm taxa both in the sandstone and lignite floral assemblages. Only such megafossils and palynomorphs which could be reliably referred to the extant taxa (genus and/or family) are considered in our survey.

### NEOGENE AND MODERN FLORISTIC COMPOSITION AND VEGETATIONAL TYPES

Almost all the information of the Neogene plant fossils (mega and micro-) of the Pondicherry-Neyveli area has been generated at Birbal Sahni Institute of Palaeobotany, Lucknow and Osmania University, Hyderabad. As of today, 53 families of Angiosperms encompassing 108 genera could be recognized reliably (Table 2). Of these, 5 (Agavaceae, Arecaceae, Araceae, Poaceae and Potamogetonaceae) represent monocots, and 48 dicots. Arecaceae, numerically is the best represented family among the monocots. *Cocos*, *Eugeissona*, *Calamus*, *Sclerosperma* and *Livistona* are the palms that could be identified. Among the dicot families Anacardiaceae, Barringtoniaceae, Clusiaceae, Combretaceae, Dipterocarpaceae and Fabaceae (Caesalpinioideae and Mimosoideae) are better represented.

C. G. K. Ramanujam

Table 2. Angiosperms from the Neogene of Pondicherry-Neyveli area

Family	Fossil wood and related modern genus	Fossil pollen and related modern genus
AGAVACEAE	Carbonized stem ( <i>Dracaena</i> )	
ALANGIACEAE	<i>Alangioxylon</i> ( <i>Alangium</i> )	<i>Lonagiopollis</i> ( <i>Alangium</i> )
ANACARDIACEAE	<i>Manigiferoxylon</i> ( <i>Mangifera</i> ) <i>Glutoxylon</i> ( <i>Gluta</i> ) <i>Bouea</i> ( <i>Bouea</i> )	<i>Rhoipites</i>
ARALIACEAE		<i>Araliaceopollenites</i> ( <i>Aralia</i> )
ARECACEAE	<i>Palmoxylon</i> ( <i>Livistona</i> )	<i>Palmaepollenites</i> ( <i>Cocos</i> ) <i>Arecipites</i> <i>Quilonipollenites</i> ( <i>Eugeissona</i> ) <i>Longapertites</i> <i>Proxapertites</i> <i>Dicolpopollis</i> ( <i>Calamus</i> ) <i>Trilatiporites</i> ( <i>Sclerosperma</i> ) <i>Spinainaperturites</i> ( <i>Typhonium</i> )
ARACEAE		<i>Retitricolporites sitholeyi</i> ( <i>Avicennia</i> )
AVICENNIACEAE		<i>Marginipollis</i> ( <i>Barringtonia</i> )
BARRINGTONIACEAE	<i>Barringtonioxylon</i> ( <i>Barringtonia</i> ) <i>Careyoxylon</i> ( <i>Careya</i> )	
BOMBACACEAE		<i>Bombacacidites</i> <i>Lakiapollis</i> ( <i>Cullenia</i> )
BORAGINACEAE	<i>Cordioxylon</i> ( <i>Cordia</i> )	
BRASSICACEAE		<i>Cruciferoipollenites</i>
CARYOPHYLLACEAE		<i>Polyporina</i>
CHENOPODIACEAE		<i>Polyporina globosa</i>
CLUSIACEAE	<i>Calophylloxylon</i> ( <i>Calophyllum</i> )	<i>Pachydermites</i> ( <i>Symphonia</i> ) <i>Pentadesmapites</i> ( <i>Pentadesma</i> )
	<i>Mesuoxydon</i> ( <i>Mesua</i> )	
COMBRETACEAE	<i>Terminalioxylon</i> ( <i>Terminalia</i> ) <i>Anogeissuoxylon</i> ( <i>Anogeissus</i> )	<i>Heterocolpites</i> ( <i>Lumnitzera</i> )

## Floristic diversity in the Neogene and Modern forests

Family	Fossil wood and related modern genus	Fossil pollen and related modern genus
CTENOLOPHONACEAE		<i>Ctenolophonidites</i> ( <i>Ctenolophon</i> )
DIPTEROCARPACEAE	<i>Dipterocarpoxyton</i> ( <i>Dipterocarpus</i> ) <i>Dryobalanoxyton</i> ( <i>Dryobalanops</i> ) <i>Anisopteroxyton</i> ( <i>Anisoptera</i> ) <i>Shoreoxyton</i> ( <i>Shorea</i> ) <i>Hopenium</i> ( <i>Hopea</i> )	<i>Retitricolpites dipterocarpodites</i> ( <i>Dipterocarpus</i> )
DROSERACEAE		<i>Droseridites</i>
EBENACEAE	<i>Ebenoxyton</i> ( <i>Diospyros-Maba</i> )	
EUPHORBIACEAE	<i>Putranjivoxyton</i> ( <i>Putranjiva</i> ) <i>Bridelioxyton</i> ( <i>Bridelia</i> )	<i>Crotonoidaepollenites</i> ( <i>Jatropha</i> ) <i>Crotonipollis</i>
FABACEAE	<i>Cynometroxylon</i> ( <i>Cynometra</i> ) <i>Milletioxylon</i> ( <i>Milletia</i> ) <i>Pterocarpoxyton</i> ( <i>Pterocarpus</i> ) <i>Pterogynoxyton</i> ( <i>Pterogyne</i> ) <i>Peltophoroxyton</i> ( <i>Peltophorum</i> ) <i>Pahudioxyton</i> ( <i>Afzelia - Intsia</i> ) <i>Tamarindoxyton</i> ( <i>Tamarindus</i> ) <i>Euacacioxyton</i> ( <i>Acacia</i> ) <i>Hopeoxyton</i> ( <i>Sindora</i> ) <i>Pericopsoxyton</i> ( <i>Pericopsis</i> ) <i>Albizinium</i> ( <i>Albizia</i> )	<i>Margocolporites</i> ( <i>Caesalpinia</i> )  <i>Trisyncolpites</i> ( <i>Poinciana, Caesalpinia</i> ) <i>Palaeocaesalpinaceaepites</i>  <i>Polyadopollenites</i> ( <i>Acacia, Albizia</i> )
HALORAGACEAE		<i>Haloragacidites</i> ( <i>Myriophyllum</i> )
HIPPOCRATEACEAE		<i>Hippocrateaceaedites</i> <i>Dakshinipollenites</i>

## C. G. K. Ramanujam

Family	Fossil wood and related modern genus	Fossil pollen and related modern genus
ICACINACEAE		<i>Icacinoipollenites</i>
LAMIACEAE		<i>Polycolpites</i>
LORANTHACEAE		<i>Gothanipollis</i>
LAURACEAE	<i>Litsea (Litsea)</i>	
LENTIBULARIACEAE		<i>Neyvelipollenites (Utricularia)</i>
LYTHRACEAE	<i>Lagerstroemioxylon (Lagerstroemia)</i>	<i>Verrutricolporites rotundiporis (Crenea)</i>
MENISPERMACEAE		<i>Assamialetes</i>
MORACEAE		<i>Triporopollenites (Artocarpus)</i>
NYMPHAEACEAE		<i>Monosulcites neyveliense (Nymphaea)</i>
ONAGRACEAE		<i>Triorites</i>
OLACACEAE		<i>Anacolosidites (Anacolosia)</i>
OLEACEAE		<i>Retitricolporites cuddalorensis (Ligustrum)</i>
POACEAE		<i>Graminidites</i>
POTAMOGETONACEAE		<i>Retipilonapites (Potamogeton)</i>
PULMBAGINACEAE		<i>Warkallipollenites (Aegialitis)</i>
		<i>Plumbaginacipites</i>
POLYGALACEAE	<i>Xanthophyllum (Xanthophyllum)</i>	<i>Polygalacidites (Polygala)</i>
POLYGONACEAE		<i>Polygonacidites (Polygonum)</i>
PROTEACEAE		<i>Proteacidites (Isopogon)</i>
ROSACEAE	<i>Parinariosxylon (Parinari)</i>	
RUBIACEAE		<i>Palaeocoprosmadites (Coprosmia)</i>
RHIZOPHORACEAE	<i>Carallioxylon (Carallia)</i>	<i>Zonocostites (Rhizophora)</i>
SAPOTACEAE	<i>Chrysophylloxylon (Chrysophyllum)</i>	<i>Sapotaceoidaepollenites</i>
SAPINDACEAE	<i>Sapindoxylon</i>	<i>Talisipites</i>
		<i>Cupanieidites (Cupania)</i>
SIMAROUBACEAE	<i>Ailanthoxylon (Ailanthus)</i>	
SONNERATIACEAE	<i>Sonneratioxylon (Sonneratia)</i>	<i>Florschuetzia (Sonneratia)</i>

## Floristic diversity in the Neogene and Modern forests

Family	Fossil wood and related modern genus	Fossil pollen and related modern genus
	<i>Dubangoxylon</i> ( <i>Dubanga</i> )	
STERCULIACEAE	<i>Sterculinium</i> ( <i>Sterculia</i> )	<i>Tricollareporites</i> ( <i>Pterospermum</i> )
SYMPLOCACEAE		<i>Symplocoipollenites</i> ( <i>Symplocos</i> )
THYMELIACEAE		<i>Thymelaepollis</i> <i>Clavaperiporites</i> ( <i>Wikstroemia</i> )
TILIACEAE	<i>Grewioxylon</i> ( <i>Grewia</i> )	<i>Intratriporopollenites</i> <i>Lacrimapollis</i> ( <i>Brownlowia</i> )
ULMACEAE	<i>Holoptelioxylon</i> ( <i>Holoptelea</i> )	

(Data collected from : Agarwal, 1988, 1990a, 1990b, 1991a, 1991b; Ambawani, 1982; Lakanpal & Awasthi, 1964; Awasthi, 1965, 1967, 1969a, 1969b, 1969c, 1970a, 1970b, 1971, 1974a, 1974b, 1975, 1977a, 1977b, 1979, 1980, 1981, 1984, 1986; Awasthi & Agarwal, 1986; Navale, 1956, 1959, 1964, 1973; Navale & Misra, 1979; Bande & Ambawani, 1982; Ramanujam, 1953a, 1953b, 1954, 1956a, 1956b, 1956c, 1958, 1960, 1961, 1966a, 1966b, 1966-67, 1968; Ramanujam & Rao, 1966, 1969; Ramanujam *et al.*, 1981; Ramanujam & Reddy, 1984; Ramanujam *et al.*, 1985; Reddy, 1981, 1989; Reddy *et al.*, 1984; Alpana Singh *et al.*, 1992).

There are a fairly good number of families which are represented by more than one fossil type i. e., silicified wood, carbonized wood, pollen, cuticle and mummified leaves. These are of considerable significance in any discussion on floristics. These families include Arecaceae, Dipterocarpaceae, Anacardiaceae, Ebenaceae, Rhizophoraceae, Clusiaceae, Combretaceae, Sonneratiaceae, Barringtoniaceae, Fabaceae, Sapotaceae, Euphorbiaceae and Thymeliaceae (Ramanujam, 1968; Awasthi, 1974b; Guleria, 1992; Srivastava, 1984; Upadhyay & Verma, 1986; Verma *et al.*, 1989; Agarwal, 1990b). The overall Neogene floristic spectrum of the Pondicherry-Neyveli area clearly signifies the occurrence of pockets of brackish water or estuarine mangrove swamps flush with the coast line and tropical wet evergreen forests away from the coastal belt (Ramanujam, 1982, 1990; Alpana Singh *et al.*, 1992). These two vegetational types reflect upon the then prevailing ecological conditions. The mangrove floral complexes could be found in the littoral regions between low and high tides at the river mouths where the soil is saline and mostly silty and clayey. *Rhizophora*, *Avicennia*, *Sonneratia*, *Barringtonia*, *Lumnitzera*, *Crenea*, *Aegialitis*, *Excoecaria* and *Brownlowia* constitute the mangrove taxa in the Neogene floras of the Pondicherry-Neyveli area.

Taxa which unequivocally point towards the prevalence of tropical wet evergreen forest, a product of warm climate and extreme humidity because of heavy precipitation all through the



**C. G. K. Ramanujam**

year include *Gluta*, *Bouea*, *Calophyllum*, *Mesua*, *Pentadesma*, *Symphonea*, *Dipterocarpus*, *Dryobalanops*, *Anisoptera*, *Hopea*, *Diospyros*, *Ctenolophon*, *Cynometra*, *Milletia*, *Artocarpus*, *Anacolosia*, *Alchornia*, *Duabanga*, *Litsea*, *Parinari*, *Cullenia*, etc. (Champion & Seth, 1968; Puri *et al.*, 1983).

The members of Arecaceae are generally coastal and occupy a position away from the storm tide dominating the sandbeach flora. The deciduous taxa, viz., *Terminalia*, *Lagerstroemia*, *Anogeissus*, *Albizia*, *Cassia*, *Acacia*, *Sterculia*, etc. (Table 2) encountered along with the wet evergreens, must have occupied the more open segments in the evergreen forests apparently imparting a semi-evergreen look to these segments. When gaps are created accidentally in the otherwise dense evergreen forests resulting in relatively drier niches the deciduous elements from nearby regions creep in and get established. The generally hardy and more adaptable nature of these deciduous taxa would enable them to coexist locally with the evergreen ones. The occurrence of hydrophytic taxa such as *Nymphaea*, *Potamogeton*, *Myriophyllum* and *Utricularia* indicate the prevalence of local ponds. The few temperate members viz., *Symplocos*, *Ligustrum* and *Wikstroemia* encountered occasionally could have had their source in some uplands around the depositional site.

In comparison to the rich Neogene vegetational complex of tropical moist evergreen type, only few discontinuous patches of natural vegetation are now encountered in the Pondicherry-Neyveli area. These patches constitute the scrub woodlands/thickets or the so-called dry evergreen forests. The hilly ranges towards the west not far from the Pondicherry environs viz., Gingee, Tiruvannamalai, Tirukkivilur, Javadi, Chengam, etc. are clothed with dry deciduous forests. Many of the deciduous members of the Neogene floras continue to be represented in these forests. Under wetter climatic regime, however, of the Neogene times, these hilly ranges could have been occupied by moist evergreen vegetation. As against the warm humid climate of the Neogene period, the present day climate of the area is considerably dry and of dissymmetric regime with the peak of rainfall during October-November months. Some rains (about 330 mm) no doubt, are received during the south-west monsoon season from June to September; but the bulk of the rain (about 800 mm), however, is during the north-east or retreating monsoon from October to January (Meher-Homji, 1974a, 1974b).

The scrub woodland formation shows taxa characteristic to it along with those seen commonly in the dry deciduous forests of the peninsula. The following are some of the more important taxa of this scrub jungle (for detailed information on this flora see Meher-Homji, 1974a; Puri *et al.*, 1983) : *Albizia amara*, *Manilkara hexandra*, *Pterospermum suberifolium*, *Atlantia monophylla*, *Garcinia spicata*, *Memecylon umbellatum*, *Maba buxifolia*, *Gmelina asiatica*, *Randia malabaricum*, *Grewia rhamnifolia*, *Sapindus emarginatus*, *Zizyphus xylopyrus*, *Chloroxylon swietinea*, *Albizia lebbek*, *Dalbergia paniculata*, *Dichrostachys cinerea*, *Flacourtia indica*, *Toddalia asiatica*, etc.

The occurrence of wet evergreen forest in the Pondicherry-Neyveli area now at a latitudinal position of about 12°N, could only be possible if this area was at the equatorial

### Floristic diversity in the Neogene and Modern forests

latitude during the Miocene age (about 20 million yrs. ago). Incidentally as visualized by Meher-Homji (1976, 1978, 1980), the evidence from the northward drift of the Indian plate at 5–6 cm per year if worked backwards would put southern India at the equatorial latitude. As India moved from an equatorial position in Miocene to its present northern latitudinal position, it experienced considerable decline in rainfall. The dramatic deterioration in the climate from Neogene to the present day could also be attributable to the final phase of the Himalayan uplift manifested during Late Pliocene through Early Pleistocene. It is during this final heave of the Himalayas that the monsoon pattern so characteristic of the Indian peninsula must have been firmly established resulting in drastic decline in the annual quantum of rainfall towards the Coromandel coast of southern India (Meher-Homji, 1980) which led to the disappearance of moist evergreen forest type leaving behind the deciduous ones. Further, the maximum elevation of the Western Ghats by the end of Pliocene effectively cut off the bulk of south-west monsoon from Coromandel coast (Meher-Homji, 1974b).

### PHYTOGEOGRAPHIC CONSIDERATIONS

Most of the Neogene rain forest taxa of the Pondicherry-Neyveli region are now encountered in Kerala state and Western Ghats where the climate continues to be of the warm humid type with heavy precipitation. It is pertinent to note that these taxa were also recorded in the Neogene (Quilon and Warkalli beds) of Kerala (Guleria, 1992). *Gluta* (Anacardiaceae), *Dipterocarpus*, *Hopea* (Dipterocarpaceae), *Cynometra* (Caesalpinioideae), *Parinari* (Rosaceae), *Alchornia* (Euphorbiaceae), *Cullenia* (Bombacaceae), *Anacolosia* (Olacaceae), etc. as evidenced from their occurrence in the Neogene floras of various parts of India especially in north-eastern region (Guleria, 1992) seem to have enjoyed a much wider distribution during the Mio-Pliocene times compared to their present day restricted spread. Further, mention may particularly be made of some members viz., *Eugeissona* (*Quilonipollenites*) and *Sclerosperma* (*Trilatiporites*) of Arecaceae, *Anisoptera* (*Anisopteroxylon*) and *Dryobalanops* (*Dryobalanoxylon*) of Dipterocarpaceae, *Pentadesma* (*Pentadesmapites*) and *Symphonia* (*Pachydermites*) of Clusiaceae, *Bouea* of Anacardiaceae, *Sindora* (*Hopeoxylon*) of Fabaceae and *Ctenolophon* (*Ctenolophonidites*) of Ctenolophonaceae, which were recorded in the Neogene strata of the Pondicherry-Neyveli (and also of Kerala) region, but are conspicuous by their absence in the modern floras of India. *Eugeissona* is a coastal palm in Southeast Asia, and *Sclerosperma* is a tropical African palm. *Anisoptera* is now confined to the wet evergreen forests of Chittagong (Bangladesh), Burma, Malayan peninsula, Sumatra, Borneo and New Guinea. *Dryobalanops* is currently restricted to the rain forests of Sumatra, Borneo and Malayan peninsula. *Bouea* is another Malayan taxon. *Pentadesma* and *Symphonia* are essentially tropical African taxa, and lastly *Ctenolophon* is now seen in West Africa, and Malayan-Phillippine area.

The Malayan elements could have entered into India after the Indian plate established connection with Asian main land which seems to have happened during the Early Miocene. This was also the time when members of Dipterocarpaceae entered into India from their original home, the Malaysia.

## C. G. K. Ramanujam

The above conspectus thus clearly highlights that the Pondicherry-Neyveli area witnessed dramatic climo-vegetational changes from wet evergreen to deciduous to scrub woodland type since Mio-Pliocene times. In conclusion, it may be said that a meaningful study of the bygone floristics and vegetational complexes would add a new dimension and a new perspective to our understanding of the modern vegetational types.

## Literature cited

- Agarwal, A. 1988. Occurrence of *Bouea* in the Neyveli lignite deposits, India. *Geophytology* 18 : 166-168.
- Agarwal, A. 1990a. *Careyxylon pondicherriense* Awasthi from Neyveli lignite deposits. *Vegetos* 3 : 179-181.
- Agarwal, A. 1990b. Leaf impressions from Neyveli lignite, Tamil Nadu, India. *J. Indian Bot. Soc.* 69 : 35-38.
- Agarwal, A. 1991a. Fossil wood of *Grewia* from Neyveli lignite deposits, India. *Vegetos* 4 : 5-7.
- Agarwal, A. 1991b. Occurrence of *Altingia* and *Bauhinia* in the Neyveli lignite (Miocene), India. *J. Indian Bot. Soc.* 70 : 119-121.
- Alpana Singh, B. K., Misra, B. D. Singh & G. K. B. Navale. 1992. The Neyveli lignite deposits (Cauvery basin), India : Organic composition, age and depositional pattern. *Intern. Jour. Coal. Geol.* 21 : 45-97.
- Ambawani, N. 1982. Occurrence of a fossil axis belonging to Agavaceae from Neyveli lignite, South India. *Geophytology* 12 : 322-324.
- Awasthi, N. 1965. Fossil woods of Anacardiaceae from the Tertiary of South India. *Palaeobotanist* 14: 131-143.
- Awasthi, N. 1967. Fossil wood resembling that of *Milletia* from the Tertiary of South India. *Curr. Sci.* 36 : 180-181.
- Awasthi, N. 1969a. A fossil wood of *Sonneratia* from the Tertiary of South India. *Palaeobotanist* 17: 254-257.
- Awasthi, N. 1969b. A new fossil wood resembling the genus *Parinarium* of the family Rosaceae from the Tertiary of South India. *Palaeobotanist* 17 : 317-321.
- Awasthi, N. 1969c. A fossil wood belonging to the family Alangiaceae from the Tertiary of South India. *Palaeobotanist* 17 : 322-325.
- Awasthi, N. 1970a. On the occurrence of two new fossil woods belonging to the family Lecythidaceae in the Tertiary rocks of South India. *Palaeobotanist* 18 : 62-74.
- Awasthi, N. 1970b. A fossil wood of Ebenaceae from the Tertiary of South India. *Palaeobotanist* 18: 192-196.
- Awasthi, N. 1971. Revision of some dipterocarpaceous woods previously described from the Tertiary of South India. *Palaeobotanist* 18 : 226-234.
- Awasthi, N. 1974a. Occurrence of some dipterocarpaceous woods in the Cuddalore series of South India. *Palaeobotanist* 21: 339-351.

## Floristic diversity in the Neogene and Modern forests

- Awasthi, N. 1974b. Neogene Angiosperm woods. In : K. R. Surange, R. N. Laxhanpal & D. C. Bhardwaj (Eds.), *Aspects and Appraisal of Indian Palaeobotany*. BSIP, Lucknow, India. pp. 341-358.
- Awasthi, N. 1975. *Milletioxylon indicum* Awasthi, a fossil wood of Leguminosae from the Cuddalore series of South India. *Palaeobotanist* 2 : 47-50.
- Awasthi, N. 1977a. On two new fossil woods resembling *Chrysophyllum* and *Holoptelea* from the Cuddalore series near Pondicherry. *Palaeobotanist* 24 : 21-25.
- Awasthi, N. 1977b. Revision of *Hopeoxylon indicum* Navale and *Shoreoxylon speciosum* Navale from the Cuddalore series near Pondicherry. *Palaeobotanist* 24 : 102-107.
- Awasthi, N. 1979. Three new leguminous woods from the Cuddalore series near Pondicherry. *Palaeobotanist* 26 : 157-166.
- Awasthi, N. 1980. Two new dipterocarpaceous woods from the Cuddalore series near Pondicherry. *Palaeobotanist*, 26 : 248-256.
- Awasthi, N. 1981. Fossil woods belonging to Sterculiaceae and Lythraceae from the Cuddalore series of Pondicherry. *Palaeobotanist* 27 : 182-189.
- Awasthi, N. 1984. Studies on some carbonised woods from the Neyveli lignite deposits, India. *Geophytology* 14 : 82-95.
- Awasthi, N. 1986. Fossil wood of *Xanthophyllum* from the Cuddalore sandstones near Pondicherry. *Palaeobotanist* 35 : 314-317.
- Awasthi, N. & A. Agarwal. 1986. A carbonised wood resembling *Parinari* from the Neyveli lignite deposits, India. *Palaeobotanist* 35 : 57-60.
- Balasundar, N. K. 1968. Tertiary deposits of Neyveli lignite field. In : *Cretaceous-Tertiary formations of South India*. *Geol. Soc. Ind. Mem.* 2 : 256-262.
- Bande, M. B. & K. Ambawani. 1982. *Sclerosperma* type of pollen grains from the Neyveli lignite of India. *Palaeobotanist* 30 : 63-67.
- Champion, H. G. & S. K. Sethi. 1968. *A revised survey of the forest types of India*. Govt. Ind. Publ., Delhi. 403pp.
- Gowrisankaran, S., P. P. Sethi, R. Hariharan & K. P. Agarwal. 1987. Lignite deposits of India – their occurrence, depositional features and characteristics. In : R. M. Singh (Ed.), *Proc. Nat. Sem. Coal Res. Ind.* (1986), Varanasi. pp. 481-553.
- Guleria, J. S. 1992. Neogene vegetation of Peninsular India. *Palaeobotanist* 40 : 255-311.
- Kailasam, L. N. 1968. Some results of geophysical exploration over the Cretaceous – Tertiary Formations of South India. *Geol. Soc. Ind. Mem.* 2 : 153-167.
- Laxhanpal, R. N. & N. Awasthi. 1964. *Mesuoxyylon arcotense* gen. et sp. nov. a fossil dicotyledonous wood from the Tertiary of South Arcot district, Madras, India. *Palaeobotanist* 12 : 260-264.
- Meher-Homji, V. M. 1974a. The natural history of Pondicherry and its environs. *Rev. Histor. de Pondicherry* 12 : 45-54.
- Meher-Homji, V. M. 1974b. On the origin of the tropical dry evergreen forest of South India. *Int. J. Ecol. Environ. Sci.* 1 : 19-39.

## C. G. K. Ramanujam

- Meher-Homji, V. M. 1976. On the rate of drift of India during the Tertiary. *Geobios* 3 : 23-24.
- Meher-Homji, V. M. 1978. Environmental implications of Tertiary plant fossils of India. *Ind. J. Bot.* 1 : 19-27.
- Meher-Homji, V. M. 1980. Bioclimatology and vegetation of Peninsular India. *20th A. C. Seward Memorial Lecture*, BSIP, Lucknow. pp. 1-27.
- Murthy, N. G. K. 1968. The Upper Cretaceous and Tertiary Formations of Pondicherry. *In: Cretaceous-Tertiary Formations of South India. Geol. Soc. Ind. Mem.* 2 : 113-119.
- Navale, G. K. B. 1956. On two new species of *Terminalioxylon* Schonfeld from the Tertiary beds of South India. *Palaeobotanist* 4 : 35-39.
- Navale, G. K. B. 1959. Occurrence of fossil *Cynometra* from the Cuddalore series near Pondicherry, India. *Palaeobotanist* 7 : 6-11.
- Navale, G. K. B. 1964. *Anogeissusoxylon indicum* gen. et. sp. nov. from Tertiary rocks near Pondicherry, India. *Palaeobotanist* 11 : 154-158.
- Navale, G. K. B. 1973. Some contribution to the Palaeobotany of Neyveli lignite, South India. *Palaeobotanist* 20 : 179-189.
- Navale, G. K. B. & B. K. Misra. 1979. Some new pollen grains from Neyveli lignite, Tamil nadu, India. *Geophytology* 8 : 226-239.
- Puri, G. S., V. M. Meher-Homji, R. K. Gupta & S. Puri. 1983. *Forest Ecology*, Vol. 1. Phytogeography and Forest Conservation. Oxford & IBH Publ. Co., New Delhi.
- Ramanathan, S. 1968. Stratigraphy of Cauvery basin with reference to its oil prospects. *In: Cretaceous-Tertiary Formations of South India. Geol. Soc. Ind. Mem.* 2 : 153-167.
- Ramanathan, S. 1979. Tertiary Formations of South India. *Proc. Colloq. Palaeontological studies in southern region. Geol. Surv. Ind. Misc. Publ.* 45 : 165-180.
- Ramanujam, C. G. K. 1953a. *Palmoxylon arcotense* sp. nov. a fossil palm resembling the living genus *Livistona* from South India. *Palaeobotanist* 2 : 89-91.
- Ramanujam, C. G. K. 1953b. On two new species of *Mesembrioxylon* from the vicinity of Pondicherry, South India. *Palaeobotanist* 2 : 101-106.
- Ramanujam, C. G. K. 1954. On some silicified woods from near Pondicherry, South India. *Palaeobotanist* 3 : 40-50.
- Ramanujam, C. G. K. 1956a. Fossil woods of Dipterocarpaceae from the Tertiary of South Arcot district, Madras. *Palaeobotanist* 4 : 45-56.
- Ramanujam, C. G. K. 1956b. On two new species of *Terminalioxylon* Schonfeld from the Tertiary of South Arcot district, Madras. *J. Indian Bot. Soc.* 35 : 103-113.
- Ramanujam, C. G. K. 1956c. Fossil woods of Euphorbiaceae from the Tertiary rocks of South Arcot district, Madras. *J. Indian Bot. Soc.* 35 : 284-307.
- Ramanujam, C. G. K. 1958. *Palmoxylon puratanam*, a new species of petrified palms from the Tertiary rocks of South Arcot district, Madras. *J. Indian Bot. Soc.* 37 : 128-137.

**Floristic diversity in the Neogene and Modern forests**

- Ramanujam, C. G. K. 1960. Silicified woods from Tertiary rocks of South India. *Palaeontographica* 106B : 99-140.
- Ramanujam, C. G. K. 1961. A fossil dicotyledonous wood resembling the modern *Tamarindus* from the Tertiary rocks of South Arcot district, Madras. *Palaeobotanist* 8 : 38-42.
- Ramanujam, C. G. K. 1966a. A further investigation of the ligneous fossils of Combretaceae from South India. *Palaeobotanist* 14 : 246-255.
- Ramanujam, C. G. K. 1966b. Palynology of the Miocene lignite from South Arcot district, Madras. *Pollen et Spores* 8 : 149-203.
- Ramanujam, C. G. K. 1966-67. Pteridophytic spores from the Miocene lignite of South Arcot district, Madras. *Palynol. Bull.* 1-2 : 29-40.
- Ramanujam, C. G. K. 1968. Some observations on the flora of the Cuddalore sandstone series. *Mem. Geol. Soc. Ind.* 2 : 271-285.
- Ramanujam, C. G. K. & M. R. R. Rao. 1966. The occurrence of *Cynometroxylon indicum* Chowdhury & Ghosh from the Cuddalore series. *Curr. Sci.* 35 : 158-159.
- Ramanujam, C. G. K. & M. R. R. Rao. 1969. *Shoreoxylon kraeuseli* sp. nov. a new dipterocarpaceous wood from the Cuddalore series of South India. In : H. Santapau, et al. (Eds.), *J. Sen. Mem. Vol., Bot. Soc. Bengal, Calcutta* : 253-258.
- Ramanujam, C. G. K. & P. R. Reddy. 1984. Palynoflora of Neyveli lignite. Floristic and palaeoenvironmental analysis. *Jour. Palynol.* 20 : 58-74.
- Ramanujam, C. G. K., P. R. Reddy & P. S. Sarma. 1985. Additions to palynoflora of the Neyveli lignite, Tamil Nadu. *Jour. Palaeont. Soc. Ind.* 30 : 49-53.
- Ramanujam, C. G. K., K. Srisailam & P. R. Reddy. 1981. The genus *Crassoretitriletes* Germeraad, Hopping & Muller 1968 from the South Indian Tertiary deposits and its stratigraphic importance. *Geoscience* II : 1-6.
- Reddy, P. R. 1981. Investigations of plant fossils from Neyveli lignite. Ph. D. Thesis, Osmania University, Hyderabad.
- Reddy, P. R. 1989. Fossil wood of Sterculiaceae from Neyveli lignite, Tamil Nadu. *Proc. Spec. Ind. Geophytol. Conf. Pune, 1986.* pp. 133-135.
- Reddy, P. R., K. Srisailam & C. G. K. Ramanujam. 1984. The genus *Trisyncolpites* Kar of Caesalpiniaceae affinities from the Neyveli lignite, Tamil Nadu. *Ind. J. Bot.* 7 : 54-55.
- Srivastava, R. K. 1984. Leaf cuticles of *Litsea* from the Tertiary lignite of Neyveli, South India. *J. Indian Bot. Soc.* 63 : 25-28.
- Subramanyan, V. 1969. Geology and ground water aspects of the Neyveli lignite field, South Arcot district, Madras State. *Mem. Geol. Surv. Ind.* 94 : 1-298.
- Upadhyay, N. & C. L. Verma. 1986. Palm cuticle from Neyveli lignite, South India. *Geophytology* 16 : 142-143.
- Verma, C. L., N. Upadhyay & R. K. Srivastava. 1989. Leaf cuticles from the Neyveli lignite of India. *Palaeobotanist* 37 : 192-198.