


Some rare grasses recollected after a lapse of hundred years from Bihar

Jha R.R. & S.K. Varma



How to cite:

Jha R.R. & S.K. Varma 1992. Some rare grasses recollected after a lapse of hundred years from Bihar. *Rheedea* 2(1): 69–72.

https://dx.doi.org/10.22244/rheedea.1992.02.01.10

Published in print: 30.06.1992 Published Online: 01.01.2022

Vol. 2 (1): 69-72 1992

Some rare grasses recollected after a lapse of hundred years from Bihar

R. R. Jha and S. K. Varma

Department of Botany, Bhagalpur University, Bhagalpur-812007.

Abstract

26 taxa of the family Poaceae have been recollected from Bihar after a lapse of about hundred years.

Poaceae is one of the largest family of Bihar and contains more than 300 species (Jain et al, 1975). During a study of the grass flora of Bihar and scanning of specimens at Central National Herbarium, Howrah, the authors could determine that the following grasses - Aristida funiculata Trin. et Rupr., Brachiaria kurzii (Hook. f.) A. camus, Chloris montana Roxb., Chrysopogon lancearius (Hook. f.) Haines, Coelachne simpliciuscula (Wight. & Arn. ex steud.) Munro ex Benth., Coix aquatica Roxb., Cynodon arcuatus J. S. Presl ex C. B. Presl, C. barberi Rang, et Tad., Dimeria ornithopoda Trin., Dinebra retroflexa (Vahl) Panz., Garnotia tenella (Arn. ex Miq.) Jan., Hymenachne pseudo-interrupta C. Muell., Ischaemum indicum (Houtt.) Merr. var. indicum subvar. malacophyllum (Hochst.) Bor, Iseilema laxum Hack., Melanocenchris jacquemontii Jaub, et Spach, Microstegium ciliatum (Trin.) A. Camus, Mnesithea laevis (Retz.) Kunth, Narenga porphyrocoma (Hance) Bor, Oropetium thomaeum (L. f.) Trin., Panicum antidotale Retz., Pseudopogonatherum contortum (Brongn.) A. Camus, Pseudoraphis brunoniana Griff., Schoenefeldia gracilis Kunth, Sporobolus tetragonus Bor, Sehima nervosum (Rottl.) Stapf and Tragus roxburghii Panigrahihave been recollected from Bihar after a lapse of about hundred years. Perusal of literature reveals that the above mentioned grasses have not been collected by Mooney (1950), Bressers (1951), Panigrahi (1966), Paul (1967), Thothathri et al. (1966), Banerjee & Banerjee (1968), Manna & Samaddar (1984) and Singh (1985). Recollection of the above rare grasses from different localities of Bihar is, hence interesting. The history of their earlier collection and the localities of the recent collection are provided here. All the cited specimens are deposited in Bhagalpur University Herbarium, Bhagalpur.

Aristida funiculata Trin. et Rupr.

J. D. Nusker collected this annual grass from Gaya (Gaya, Oct. 1902, Nusker s. n. CAL acc. no 533997). We have been able to collect it from Gilamari, 3km south-east from Mirzachowki, Sahibganj District. (RRJha 6493).

Brachiaria kurzii (Hook. f.) A. Camus.

This rare grass was once collected

from Bihar (Parasnath; Hooker 12, CAL). It has been recollected from Mirzachowki Sahibganj District. (RRJha 6576).

Chloris montana Roxb.

Mokim collected this rare grass from Rajmahal of Bihar during Oct. 1894, (Mokim 1464, CAL). Now it has been collected from the dry sandy soils of Gandhigram (Godda District) (RRJha 6532) and Shripur, 18 km south from Mirzachowki, Sahibganj District (RRJha 6534).

Chrysopogon lancearius (Hook. f.) Haines

This perennial grass, which grows in rock crevices, was earlier collected by Kurz, Clarke and Haines from Bihar (Munger hills, 3. 11. 1868, Kurz s. n. CAL acc. no. 531018; Ramghur ghat, Sing-bhum, 1750°, 19. 10. 1883, Clarke 33898 B, CAL; Singbhum, Sept. 1899, Haines 201, CAL). We have recollected the same from Mirzachowki, Sahibganj District, (RRJha 6573) and Gandhigram, Godda District. (RRJha 7805 A.).

Coelachne simpliciuscula (wight. & Arn. ex Steud.) Munro ex Benth.

Earlier this grass was collected from Ranchi District of Bihar (Chotanagpur, Ranchi, 2000', 4. 12. 1874, *Clarke* 25190, CAL). The same has been recollected from Dumka District. Mahuagarhi, 1800', (*RRJha* 6506).

Coix aquatica Roxb.

Kurz and Mokim had collected this rare grass from Bihar (Purneah, 30. 10. 1868, Kurz s. n. CAL. acc. no. 525525; Teenpahar, 2. 11. 1868; Kurz 199, CAL; Munger, Oct. 1894, Mokim 1397, CAL). The same has been recollected by us from Simartala, Sahibganj District, (RRJha 6345).

Cynodon arcuatus J. S. Presl ex C. B. Presl

Earlier, it has been collected by Kurz and Nusker from Bihar (Baidyanath, 7. 10. 1894, *Nusker* s. n.; Rajmahal Hills, Sahibganj Oct. 1870; *Kurz* 714, CAL), We have collected it from Khutnashi. Bhagalpur District. (*RRJha* 6733); Karamtola, Sahibganj District, (*RRJha* 6443) and Sundarpahari, Godda District, (*RRJha* 6529).

Cynodon barberi Rang. et Tad.

Previously, the plant was collected by Clarke and Nusker (Hazaribag, 12. 10. 1883, *Clarke* 33784 B, CAL; Baidyanath, 10. 10. 1984, *Nusker* 1279, CAL). Recently, the same has been collected from Shripur, Sahibganj Disirict, (*RRJha* 6553),

Dimeria ornithopoda Trin.

This rare grass was first collected by Clarke from Singbhum District of Bihar (Cheybasa, 1500', 20. 11. 1883, Clarke 34350 A, CAL). The same has been recollected from Maharajpur, Sahibganj District, (RRJha 7198).

Dinebra retroflexa (Vahl) Panz.

Kurz collected this rare grass from Bihar without mentioning any specific locality (Behar, Kurz s n., CAL, acc. no. 539424). The same has been recollected from Karamtola, Sahibganj District, (RRJha 6550).

Garnotia tenella (Arn ex Miq.) Jan.

The only earlier collection is from Parasnath by Clarke (Parasnath, Hazaribag, 4300' 10. 10. 1873, *Clarke* 21057 E; 7. 10. 1883, *Clarke* 33091 C, CAL). The same has been collected on wet moss

covered rocks at Motijharna, Maharajpur, Sahibganj District, (*RRJha* 6363).

Hymenachne pseudo-interrupta C. Muell.

This rare aquatic grass was earlier recorded from Santhal Pargana by Mokim (Rajmahal, Oct. 1894, *Mokim* 1474, CAL). The same has been collected along the muddy bank of tributary of river Ganga at Babupur, Sahibganj District (*RRJha* 6428) and Hazipur diara, Bhagalpur District, (*RRJha* 6612).

Ischaemum indicum (Houtt.) Merr. var. **indicum** subvar. **malacophyllum** (Hochst.) Bor.

This prostrate annual grass was first reported from Bihar by Clarke and Kurz (Hazaribag, 10. 10. 1873, *Clarke* 21187 B, CAL; "Behar", *Kurz* 452, CAL). It has been recollected from Karamtola of Sahibganj District (*RRJha* 6609).

Iseilema laxum Hack.

This marshy grass was earlier collected from Bihar by Clarke (Gaya, 10. 10. 1894, *Mokim* 1354, 1357, 1358, 1377, CAL). We have collected it from Asanbani, Dumka District, (*NNDas* 2939) and Mirzachowki, Sahibganj District, (*RRJha* 6586).

Melanocenchris jacquemontii Jaub. & Spach

The only earlier collection is from Gaya by Mokim (Gaya, Oct. 1894, *Mokim* 1368, CAL). The same has been collected from another locality, Gandhigram in Godda District, (*RRJha* 7091).

Microstegium ciliatum (Trin.) A. Camus

This grass was collected by Clarke from two places of Bihar (Hazaribag, 10. 10. 1883. *Clarke* 33794 B; Parasnath, *Clarke* 24859, CAL). We could collect this rare grass from Maharajpur, Sahibganj District, (*RRJha* 6604), Chunjo hill, Dumka District, (*RRJha* 6629) and Angwali, Godda District, (*RRJha* 7347).

Mnesithea leavis (Retz.) Kunth

The lone collection of this grass from Bihar is of Mokim (Munger, Oct. 1894, Mokim 1405, CAL). This grass is now fairly common and it has been collected from Shivnarayanpur, Bhagalpur District (RRJha 5830), Mirzachowki, Sahibganj District, (RRJha 6636) and Guljarbag, Godda District, (RRJha 6640).

Narenga porphyrocoma (Hance) Bor

This perennial grass was earliar collected from Bihar by Kurz, Clarke and Haines (Purneah, 30. 10. 1868, Kurz s. n., CAL acc. no. 527469; Parasnath, 4000', 10. 10. 1873, Clarke 21085 G, CAL; Singbhum, Dec. 1902, Haines 494, CAL). The same has been collected from Shivnarayanpur, Bhagalpur District, (RRJha 5841), Narpatganji, Saharsa District, AKSinha 705; 714) and Karamtola, Sahibganj District, (RRJha 6608, 6617).

Oropetium thomaeum (L. f.) Trin.

Mokim was the first person who collected this annual grass from Bihar (Gaya, Oct. 1894, *Mokim* 1349; Munger, *Mokim* 1416, CAL). This rare grass has now been recollected from Jagarnathpur, Godda District, (*RRJha* 7011).

Panicum antidotale Retz.

Clarke was the only person who collected this perennial, waste-land grass from Bihar (Parasnath, 4000', 2. 10. 1873, *Clarke* 20779 A, CAL). Recently the same was collected from Sahibganj District, Mirzachowki (*RRJha* 7375).

Pseudopogonatherum contortum (Brongn.) A. Camus.

This annual grass was collected by Clarke and Gamble from Bihar (Bundgao, 2000', Singbhum district, 28. 10. 1873, Clarke 33835, CAL; Palamu, Dec. 1880, Gamble 8789, CAL). This has been recollected from the mixed forest of Kathi-

kund, Dumka District (*NNDas* 3927, *RRJha* 6616) and Angwali. Godda District, *RRJha* 6722).

Pseudoraphis brunoniana Griff.

This floating grass was first collected by Mokim from Bihar (Gaya, Oct. 1894, Mokim 1334, CAL), but it has not been included in *The Botany of Bihar and Orissa* by Haines (1925). The same has been recollected from a tank at Shibpur, Godda District, (RRJha 7071).

Schoenefedia gracilis Kunth

Inclusion of this grass in Haines, Flora is solely based on a collection of Anderson (Haines, 1925). We could collect it from Babupur diara, Sahibganj District, (*RRJha* 6781).

Sporobolus tetragonus Bor

The only collection of this annual grass from Bihar was made by Clarke (Rampur ghat, Chotanagpur, 1500'. 19. 10. 1883, *Clarke* s. n., CAL). Now we could

collect this rare grass from Angwali, Godda District, (RRJha 6785).

Sehima nervosum (Rottl) Stapf

C. B. Clarke collected this grass from Ranchi and Hazaribag districts of Bihar (Hazaribag, 1750', 13. 10. 1883, *Clarke* 35791 B; Ranchi, 2000', 16. 10. 1873, *Clarke* 21226 B, CAL). The same has been recollected along stream banks near Karamtola, Sahibganj District (*RRJha* 5862) and Angwali, Godda District (*RRJha* & *SKVarma* 7290).

Tragus roxburghii Panigrahi

This curious grass was earlier collected from Singbhum by Haines (1925). The authors were able to collect this litile grass from Bharjor, Dumka district, (NNDas 3636), Kathikund, Dumka district, (RRJha 6352), Angwali, Godda district, (RRJha 7374) and Mirzachowki, Sahibganj, District, (RRJha 6553).

Acknowledgements

Authors are grateful to Prof. K. S. Bilgrami, Head, Department of Botany, Bhagalpur University, for providing necessary facilities. We are also thankful to the Joint-Director, Central National Herbarium, Howrah, for permission to consult the Herbarium, and to CSIR, New Delhi for financial assistance.

Literature cited

- Banerjee, S. P. & Banerjee, R. N. 1968. Materials for a Flora of North Bihar-1. Champaran District. *Bull. Bot. Surv. India* 10: 183-189.
- Bressers, J. 1951. The Botany of Ranchi District, Bihar, India. Ranchi.
- Haines, H. H. 1925. The Botany of Bihar and Orissa, London.
- Jain, S. K., Banerjee, D. K. & Pal, D. C. 1975. Grasses of Bihar, Orissa and West Bengal. J. Bombay nat. Hist. Soc. 72 (3): 758-773.
- Manna, M. K. & Samaddar, U. P. 1984. On a Collection of plants from Netarhat plateaux. Palamau District, Bihar - A critical area of botanical importance. Bull. Bot. Surv. India. 26:133-141.

- Mooney, H. F. 1950. Supplement to the Botany of Bihar and Orissa. Ranchi.
- Panigrahi, G. 1966. A botanical tour in the Rajmahał Hills of Bihar. *Bull Bot*, *Surv. India*. 8:1-15.
- Paul, S. R. 1967. Studies on the grasses of Bhagalpur District (Bihar). *Ind. For.* 93:169-
- Singh, K. L. B. 1985. Grasses of Saharsa District, Bihar, J. Econ. Tax. Bot. 6(1): 245-248.
- Thothathri, K., Shetty, B. V. & Hazara P. K. 1966. A contribution to the flora of Udaipur forests in Champaran District, North Bihar. Bull. Bot. Surv. India. 8: 133-141.