




## *Teramnus hookerianus*, a heterotypic synonym of *T. labialis* (Leguminosae: Papilionoideae)

Vinay Ranjan



How to cite:

**Ranjan V. 2016.** *Teramnus hookerianus*, a heterotypic synonym of *T. labialis* (Leguminosae: Papilionoideae). *Rheedia* 26(1): 64-66.

<https://dx.doi.org/10.22244/rheedia.2016.26.01.11>

Received: 23.7.2015

Revised and accepted: 22.5.2016

Published in print: 30.06.2016

Published Online: 30.06.2016



Published by Indian Association for Angiosperm Taxonomy

This volume of Rheedia is published with partial financial assistance from Science Engineering Research Board, Government of India, New Delhi

# *Teramnus hookerianus*, a heterotypic synonym of *T. labialis* (Leguminosae: Papilionoideae)

Vinay Ranjan

Central National Herbarium, Botanical Survey of India, PO Botanic Garden, Howrah – 711 103, West Bengal, India.  
E-mail: ranjanv\_alld@rediffmail.com

## Abstract

*Teramnus hookerianus* A. Sen is relegated here to a heterotypic synonym of *T. labialis* (L.f.) Spreng.

**Keywords:** Heterotypic Synonym, Papilionoideae, *Teramnus hookerianus*, *T. labialis*

## Introduction

The genus *Teramnus* P. Browne (Leguminosae: Papilionoideae) is represented by nine species and is distributed in tropical regions of the world (Lewis *et al.*, 2005). In India, the genus is represented by four species and one subspecies (Sanjappa, 1992). During the taxonomic study of *Teramnus* for flora of West Bengal, it was observed that the species *T. hookerianus* A. Sen is represented only by the type specimens and its diagnostic characters are overlapping with another widespread species, *T. labialis* (L.f.) Spreng.

Sen (1976) described *T. hookerianus* based on the collections from Manbhum in West Bengal and Madras Presidency (without precise locality). He distinguished his new species from *T. labialis* in leaves being glabrous versus adpressed-hairy above and glabrous beneath, leaflets smaller (1–1.9 cm long) versus larger (1–5 cm long), stipules smaller (c. 4.5 mm long) versus larger (4–6 mm long) and fruits smaller (35–40 × c. 2.5 mm) versus larger (32–65 × 2.3–4 mm). However, an examination of type specimens shows that these diagnostic characters are not stable and are very variable even in the holotype (*V. Ball* s.n.) and paratypes (*M.S. Ramaswami* 1015 and *M.N. Sanyal* 532) housed at CAL (**Fig. 1b & c**). Therefore, to support the observation, specimens of *T. labialis* at CAL (**Fig. 1d & e**; Appendix 1) have been studied critically in addition to the lectotype specimen (LINN-HL901.15). The result of this study shows explicitly that the diagnostic characters of *T. hookerianus* fall very well within the circumscription of a widespread species, *T. labialis* and also corroborates the views of Verdcourt (1970). Therefore, *T. hookerianus*

is reduced here to a heterotypic synonym of *T. labialis*.

**Teramnus labialis** (L.f.) Spreng., Syst. Veg. 3: 235. 1826; Baker in Hook.f., Fl. Brit. India 2: 184. 1876; Prain, Bengal Pl. 1: 393. 1903; Sanjappa, Legumes India: 261. 1992. *Glycine labialis* L.f., Suppl. Pl.: 325. 1782.

Lectotype: Sweden, Uppsala, s. coll. s.n. (LINN-HL901.15, image!; designated by Verdcourt, 1970).

*Teramnus hookerianus* A. Sen, J. Bombay Nat. Hist. Soc. 73: 187. 1976, **syn. nov.**

Type: INDIA, West Bengal, Purulia district, Manbhum, *V. Ball* s.n. (Holotype, CAL0000012126!).

**Fig. 1a**

*Distribution:* Throughout India.

## Acknowledgements

Author is thankful to Dr. Paramjit Singh, Director, Botanical Survey of India, Kolkata, and to the Head, Central National Herbarium, Howrah, for providing facilities. He is also deeply grateful to Dr. S.V. Pradeep, Associate Professor, Department of Botany, S.V.R.N.S.S. College, Vazhoor, Kottayam, for valuable suggestion.

## Literature Cited

- Lewis, G., Schrire, B., Mackinder, B. & M. Lock 2005. *Legumes of the World*. Royal Botanic Gardens, Kew.
- Sanjappa, M. 1992. *Legumes of India*. Bishen Singh Mahendra Pal Singh, Dehra Dun.



Fig. 1. *Teramnus hookerianus* A. Sen: a. Holotype (V. Ball s.n., CAL0000012126); b & c. Paratypes (M.S. Ramaswami 1015, CAL0000012123 & M.N. Sanyal 532, CAL0000012124); d & e. Specimens of *T. labialis* (L.f.) Spreng. (C.B. Clarke 31813C & C.B. Clarke 31869A, CAL).

**Sen, A. 1976.** A new species of *Teramnus* Sw. (Fabaceae) from Manbhum (India). *J. Bombay Nat. Hist. Soc.* **73**: 187–188.

**Verdcourt, B. 1970.** Studies in the Leguminosae–Papilionoideae for the ‘Flora of Tropical East Africa’: II. *Kew Bull.* **24**: 235–307.

#### Appendix 1: Specimens examined

**Teramnus hookerianus:** INDIA, Madras Presidency, *s.loc.*, *s.d.*, *M.S. Ramaswami* 1015 (CAL). **West Bengal**, Manbhum, 1866–67, *V. Ball* s.n. (CAL); Bankura district, *s.loc.*, 9.2.1965, *M.N. Sanyal* 532 (CAL).

**Teramnus labialis:** INDIA, **Andhra Pradesh**, East Godavari, 16.2.1947, *V. Narayanaswami* 375 (CAL); Kurnool, 11.12.1976, *L.J.G. van der Maesen* 3955. **Assam**, *s. loc.*, *S.N. Bal* 836

(CAL). **Goa**, Angidev, 24.4.1966, *John Cherian* 109104 (CAL). **Gujarat**, Baroda district, *s. loc.*, 6.11.1982, *M. Sanjappa* 9525 (CAL). **Jharkhand**, Santhal Pargana, Trikut hill, 22.9.1992, *J.K. Das* 22301; Rajgir WLS, 12.9.2003, *P. Pillai* 32813 (CAL). **Kerala**, Travancore, *M.A. Lawson* 185; Palakkad district, Perambicolum, 3–4000 ft, November 1910, *A. Meebold* 12461 (CAL). **Maharashtra**, Thane, 25.10.1967, *K.V. Billore* 113241; Poona-Sanswad, 2.10.1957, *S.P. Mahajan* 26905 (CAL). **Odisha**, Puri, 2.12.1965, *T.A. Rao* s.n. (CAL). **West Bengal**, Malda, 18.11.1965, *R.M. Dutta* 27; Howrah, IBG, 22.11.1965, *V.S. Sharma* S-38; 24-Parganas, 28.3.1966, *R.L. Mitra* 525 (CAL).

---

Received: 23.7.2015

Revised and Accepted: 22.5.2016