


Brachiaria villosa var. *barbata* Bor (Poaceae) -
a new record for India

Salunkhe C.B., Potdar G.G. & S. R. Yadav



How to cite:

Salunkhe C.B., Potdar G.G. & S. R. Yadav 2002. *Brachiaria villosa* var. *barbata* Bor (Poaceae) - a new record for India. *Rheedea* 12(2): 175-177.

<https://dx.doi.org/10.22244/rheedea.2002.12.02.12>

Published in print: 31.12.2002

Published Online: 01.01.2022

***Brachiaria villosa* var. *barbata* Bor (Poaceae) – a new record for India**

C. B. Salunkhe, G. G. Potdar* and S. R. Yadav**

Department of Botany, Krishna Mahavidyalaya, Rethare Bk. – 415 108,
Maharashtra, India.

*Shivaji University, Kolhapur – 416 004, Maharashtra, India

**Department of Botany, Delhi University, Delhi – 110 007, India.

Abstract

Brachiaria villosa var. *barbata* Bor (Poaceae) is reported for the first time from India. A detailed description and illustrations are provided.

INTRODUCTION

During the survey of grasses of Maharashtra State, the authors collected an interesting species of *Brachiaria* from Devdari hills in Satara district and identified it as *Brachiaria villosa* var. *barbata*, an endemic of Nepal.

The genus *Brachiaria* Griseb. has about 100 species (Clayton & Renvoize, 1989; Watson & Dallwitz, 1994) distributed in tropics of the Old World. It has 18 species in India (Moulik, 1997). *Brachiaria villosa* var. *barbata* Bor is so far known only from Nepal (Bor, 1960). As this species has not been recorded from India, a detailed description and illustration is provided to facilitate easy identification.

***Brachiaria villosa* (Lamk.) A. Camus in Lecomte, Fl. Gen. de l'Indochine 7: 433. 1922, var. *barbata* Bor, Grass. Burma, Ceylon, India & Pakistan 286. 1960 (Fig. 1).**

Annuals. Culms erect or geniculate, 9-25 cm tall; nodes bearded. Leaves: sheaths densely hirsute with tuberculate hairs, 1.2-3.7 cm long; ligule ciliate; blades ovate-oblong, 0.8-4 x 0.3-1.2 cm, densely hirsute on both surfaces, margins undulate, scaberulous, apex subacute. Racemes 2-7, erect or ascending, 0.3-1.2 cm long; rachis slender, triquetrous, flexuous, hairy. Pedicels short, with long hairs. Spikelets broadly ovate-oblong, 2-2.5 mm long. Lower glume thinly membranous, broadly ovate-elliptic, 0.7-1 mm long, obscurely nerved, hispid on the back, apex rounded. Upper glume membranous, elliptic, 1.2-1.8 mm long, gibbous, 3-5-nerved, hispid, with subapical fringe of bristle like hairs on the back, apex mucronulate. Lower lemma membranous, ovate-oblong, 1.5-2 mm long, 3-5-nerved, hispid, with subapical fringe


C.B. Salunkhe *et al.*

Fig. 1. *Brachiaria villosa* var. *barbata* Bor: a. Habit; b. Ligule; c. & d. Spikelets; e. Lower glume; f. Upper glume; g. Lower lemma; h. Palea; i. Upper lemma; j. Palea; k. Stamens & pistil; l. Lodicules; m. & n. Caryopsis.

***Brachiaria villosa* var. *barbata* Bor. (Poaceae) – a new record**

of bristle-like hairs on the back, 2-keeled, apex mucronate. Palea hyaline, oblong, 1.4-1.8 mm long, 2-keeled, margins inflexed, apex truncate. Upper lemma coriaceous, broadly elliptic-oblong, 1.4-1.8 mm long, 5-nerved, glabrous, minutely rugulose, apex acute. Palea chartaceous, broadly elliptic-oblong, 1.4-1.8 mm long, margins inflexed, apex acute. Lodicules 2. Stamens 3; anthers 0.8-1 mm long. Ovary linear, 0.5 mm long. Grain globose, 1.1 x 0.8 mm.

Flowering and fruiting: July-September.

Distribution: So far reported only from Nepal and the present collection is a record of its extended distribution.

Specimen examined: INDIA, Maharashtra, Satara district, Devdari, 25 Aug 2000, Salunkhe 8991 (SUK).

Acknowledgements

The authors are thankful to Dr. T.A. Cope, Royal Botanic Gardens, Kew for confirming the identity of the species, and to the Ministry of Environment & Forests, Government of India for financial assistance.

Literature cited

- Bor, N. L. 1960. *The Grasses of Burma, Ceylon, India and Pakistan*. Pergamon Press, Oxford. pp.278-286.
- Clayton, W. D. & S. A. Renvoize. 1989. *Genera Graminum. Grasses of the World. Kew Bull. Addit. Ser. 13*. pp.283-284.
- Moulik, S. 1997. *The Grasses and Bamboos of India*. Scientific Publishers, Jodhpur. pp.74-80.
- Watson, L. & M. J. Dallwitz. 1994. *The Grass Genera of the World*. (Revised ed.) CAB International, U.K. pp.179-180.