

Ficus geocarpa (Moraceae): A new record for India from Arunachal Pradesh

Buragohain R., Gajurel P.R., Rethy P. & B. Singh



How to cite:

Buragohain R., Gajurel P.R., Rethy P. & B. Singh 2014. *Ficus geocarpa* (Moraceae): A new record for India from Arunachal Pradesh. *Rheedea* 24(2): 110-112.

https://dx.doi.org/10.22244/rheedea.2014.24.02.05

Received: 11.10.2013

Published in print: 31.12.2014

Revised and accepted: 03.09.2014 Published Online: 31.12.2014



Published by Indian Association for Angiosperm Taxonomy This volume of Rheedea is published with partial financial assistance from Department of Science and Technology, Government of India, New Delhi



Ficus geocarpa (Moraceae): A new record for India from Arunachal Pradesh

R. Buragohain, P.R. Gajurel*, P. Rethy and B. Singh

Forest Systematic & Ethnobiology Laboratory, Department of Forestry North Eastern Regional Institute of Sciences and Technology, (Deemed University) Nijuli – 791 109, Arunachal Pradesh *E-mail: prgajurel@gmail.com

Abstract

Ficus geocarpa Teijsm. *ex* Miq., so far known from Malesia, is reported here as a new record for India from Papumpare District, Arunachal Pradesh. Detailed description of the species with photographs are provided here to authenticate the record.

Keywords: Arunachal Pradesh, Ficus geocarpa, India, new record

Introduction

Ficus is one of the largest genera of angiosperms, with about 750 species of terrestrial trees, shrubs, hemi-epiphytes, climbers and creepers occurring in the tropics and subtropics of the world (Berg & Corner, 2005; Ronsted et al., 2008; Frodin, 2004). Ficus is represented by 115 taxa in India, with the maximum species richness (c. 61 spp.) in the Northeast India (Chaudhary et al., 2012). While working on floristic diversity of Moraceae in the eastern Himalayan state of Arunachal Pradesh, authors have collected more than 40 species of *Ficus*. One of the specimens collected from Doimukh area in Papumpare district found to be rare and quite distinct from the other Indian species. On examining this specimen, using the publications of King (1887-1888), Berg (2004) and Berg & Corner (2005) it is identified as Ficus geocarpa Teijsm. ex Miq. belonging to the subgenus Sycomorus and the section Sycocarpus. Subsequent thorough scrutiny of relevant literature (Berg & Nannapat, 2011; Chaudhary et al., 2012; Corner, 1965; Giri et al., 2008; Grierson & Long 1983; Haridasan & Rao 1987; Kanjilal et al., 1940; King, 1888; Wu et al., 2003) and herbarium specimens at CAL, ASSAM, ARUN and State Forest Research Institute, Itanagar (APFH) revealed that this species was not recorded from India so far. F. geocarpa was originally described from Malesia and till present known only from that country. Therefore this paper reports the first authentic record of this species from India and is an addition to the Indian flora. The species is described in detail with a color photoplate to facilitate easy identification.

Ficus geocarpa Teijsm. *ex* Miq., Ann. Mus. Bot. Lugduno-Batavi 3: 231. 1867; King G., Ann. Roy. Bot. Gard., Calcutta 1:102 t.135.1887; Corner, Gard. Bull. Singapore 21 (1): 91. 1965; Berg & Corner, Fl. Males., Ser. 1, Spermat. 17(2): 423. 2005. **Fig. 1**

Trees, up to 10 m tall, dioecious, evergreen; branchlets pubescent; trichomes 3-6 mm thick, soft; internodes hollow; bark smooth, pale brown. Leaves simple, distichous, slightly asymmetric, oblong to sub-ovate, 15-36 × 5-13 cm, rounded to sub-cordate at the narrow side and cordate at the broad side at base, denticulate to dentate at margin, acuminate at apex; strigose on the upper side, brown or brownish hirsute on the lower side; longer stiff hairs intermixed with shorter and softer white hairs; midrib hirtellous; cystoliths only beneath; basal nerves 3-5 pairs, prominent; wax glands present in the axils of a few middle and basal lateral nerves of the lamina; lateral nerves 6–10 (–12) pairs; tertiary nerves faintly reticulate on upper part of the lamina; petiole 0.5-2 cm long, brown- hirsute, longer stiff hairs intermixed with shorter and softer white hairs; stipules 2-4 cm long, white puberulous to subsericeous, partly brown (on the keel), strigose to hirsute, caducous. Figs flagelliflorous, branches hispid 1.5–3 m long, internodes 4–5 cm long, arising from the base of the trunk. Receptacles paired, pyriform, depressedglobose to turbinate, 1.5-1.8 cm across, covered with dense brown tomentum, ribbed towards the ostiole, apex concave, brownish- red to yellow at maturity; fig wall 3-4 mm thick; peduncle stout,



Fig. 1. *Ficus geocarpa* Teijsm. *ex* Miq. **a.** Habit; **b.** Leafy twig with adaxial and abaxial surfaces; **c.** Stipule; **d.** Fig bearing branch; **e.** Receptacle; **f.** Receptacle (Front view); **g.** Receptacle (L.S); **h-i.** Female flowers.

pubescent, $0.7-1.2 \times 0.2-3$ cm, peduncular bracts 3–4, caducous; basal bracts 2–3, 2–3 mm long; ostiole *c*. 2 mm across; ostiolar bracts 5–6, reddish pink; 10–15 whitish dots near the ostiole and at base of the fig; interfloral bracts absent. Fertile female flowers without perianth; style twice as long as the ovary, 1.5–2 mm long, lateral; stigma clavate. Mature achenes rhomboid, minutely tuberculate.

Flowering & Fruiting: August – January.

Distribution: Western and Central Malesia (Celebes, Java), India (Arunachal Pradesh).

Ecology: Found growing in the moist deciduous tropical forest in a shaded area on the hill slope at *c*. 150 m above sea level. Only a few individuals are noticed in the population, confined to a small area near road side at Doimukh of Papumpare district.

Specimen examined: INDIA, Arunachal Pradesh, Papumpare district, Doimukh at 200 m, 25.08.2012, *Rubul Buragohain* 824 (ARUN).

Notes: As the occurrence of the species is found in highly disturbed area and the only known population is with few individuals, the species could fall under threatened category. Further explorations may throw more light on the threat status. This species deserves special conservation efforts and the authors have initiated the vegetative propagation of the species in the NERIST campus, Nijuli, Arunachal Pradesh.

Acknowledgements

The authors are thankful to the Director, Botanical Survey of India, for providing access to library and herbarium facilities at CAL, ASSAM and ARUN. Expert comments and help received from C.C. Berg before his demise helped the authors in the characterization and identification of the *Ficus* species. The authors are grateful to the Director of NERIST for providing necessary facilities. Authors are also thankful to the reviewer for his valuable comments.

Literature Cited

- Berg, C.C. 2004. Flora Malesiana precursor for the treatment of Moraceae 6: *Ficus* subgenus *Sycomorus*. *Blumea*, **49**: 155-200.
- Berg, C.C. & E.J.H. Corner 2005. Moraceae *Ficus - Flora Malesiana*. Ser. I. 17(2). National Herbarium Nederland, RA Leiden. pp. 1-730.

Berg, C.C. & P. Nannapat 2011. Moraceae, In: Berg

CC & Nannapat P (ed.), *Flora of Thailand*. Vol. **10(4)** (Cecropiaceae & Moraceae), The Forest Herbarium, Bangkok, Thailand, pp. 475-675.

- Chaudhary, L.B., Sudhakar, J.V., Kumar, A., Bajpai, O., Tiwari, R. & G.V.S Murthy 2012. Synopsis of the Genus *Ficus* L. (Moraceae) in India. *Taiwania*, **57(2):** 193-216.
- **Corner E.J.H. 1965.** Check List of *Ficus* in Asia and Australasia with keys to identification. *Garden Bulletin Singapore*, **21**: 1-186.
- Frodin, D.G. 2004. History and concepts of big plant genera. *Taxon*, 53: 753-776.
- Giri, G.S., Pramanik, A. & H.J. Chowdhery 2008. Material for the Flora of Arunachal Pradesh, Vol.
 2 (Asteraceae – Ceratophyllaceae), Botanical Survey of India, Kolkata, pp. 388-407.
- Grierson, A.J.C. & D.G. Long 1983. Flora of Bhutan: Including a Record of Plants from Sikkim, Vol. 1 (1), Edinburgh: Royal Botanic Garden Edinburgh and Royal Government of Bhutan, pp. 90-99.
- Haridasan, K. & R.R. Rao 1987. Forest Flora of Meghalaya, Vol. 2 Bishen Singh Mahendra Pal Singh, Dehradun, pp. 823-832.
- Kanjilal, U.N., Kanjilal, P.C., De, R.N. & A. Das 1940. Flora of Assam, Vol. 4 Government of Assam, pp. 261-267.
- King, G. 1888. *Ficus* L. In: Hooker, J.D. *The Flora of British India*, Vol 5. L. Reeve and Co., London, UK. pp. 494-537.
- King, G. 1887 1888. The species of *Ficus* of the Indo-Malayan and Chinese Countries. *Ann. Roy. Bot. Gard., Calcutta* 1: 1-185.
- Ronsted, N., Weiblen, G.D., Savolainen, V. & J.M. Cook 2008. Phylogeny, biogeography, and ecology of *Ficus* section *Malvanthera* (Moraceae). *Molecular Phylogenetics and Evolution*, **48**: 12-22.
- Wu, Z., Zhou Z.K. & M.G. Gilbert 2003. Moraceae. In: Wu. Z., Raven, P.H. & D.Y. Hong (eds.), *Flora of China*, Vol 5. Science Press, Beijing, and Missouri Botanical Garden Press, St. Louis, pp. 21-73.

Received: 11.10.2013 Revised and Accepted: 03.09.2014