


A new species of *Calamus* (Arecaceae) from Silent Valley, Kerala, India

Renuka C., Sasidharan N. & P.V. Anto



How to cite:

Renuka C., Sasidharan N. & P.V. Anto 1997. A new species of *Calamus* (Arecaceae) from Silent Valley, Kerala, India. *Rheedea* 7(2): 69–71.

https://dx.doi.org/10.22244/rheedea.1997.07.02.01

Published in print: 31.12.1997

Published Online: 01.01.2022

ि विज्ञान एवं प्रौद्योगिकी विभाग DEPARTMENT OF SCIENCE & TECHNOLOGY **Published by Indian Association for Angiosperm taxonomy** This volume of Rheedea is published with partial financial assistance from Department of Science and Technology, Government of India, New Delhi

Rheedea

A new species of *Calamus* (Arecaceae) from Silent Valley, Kerala, India

C. Renuka, N. Sasidharan and P.V. Anto Kerala Forest Research Institute Peechi - 680 653, Trichur, Kerala, India.

Abstract

A new species of *Calamus* is described from Silent Valley National Park, Kerala, India.

INTRODUCTION


Evergreen and semievergreen forests of Western Ghats form the largest natural home of rattans in South India. Altogether 19 species of the genus *Calamus* are reported from this region (Fischer, 1931; Fernandez & Dey, 1970; Renuka, 1992). Recently another taxon of the same genus could be collected from Silent Valley National Park in Palakkad District of Kerala State. Closer scrutiny revealed that this is a new species and hence is described and illustrated here.

Calamus neelagiricus Renuka, sp. nov. (Fig. 1)

Palmijuncus solitarius. Vagina viridis spinis validis basibus bulbosis and 1 cm longis armata, rhachillae valde recurvatae circinatae. Fructus oblati 1 cm longi 0.5 cm lati, periantho pedicellato, squamis in seriebus 21 stramineis in medio canaliculatis.

Types: India, Kerala, Silent valley, evergreen forest, 1100 m, 10 March 1997. Anto 7100 - (fr.,) (Holotype - KFRI); Ibid., Sasidharan & Anto 3698 - fruit (Paratype - KFRI).

Solitary, moderate sized rattan. Stem reaching upto 25 m, with sheaths 2.5-5 cm diam., without sheaths 3 cm, internodes to 21 cm long. Sheath green, spiny; spines 1 cm long, sturdy, with a blubous base; knee present; ocrea inconspicuous; flagellum 3-4 m long. Leaf ecirrate, to 2 cm long; petiole to 25 cm, petiole and rachis armed on the dorsal side with spines; spines 0.5 cm long, sturdy, bulbous based, ventral side spiny at the point of attachment of leaflet with the rachis; leaflets long acuminate, regular, 46 x 2 cm, midvein ciliate dorsally, lateral veins ciliate ventrally, cilia to 0.3 cm, apical leaflets free. Inflorescence flagellate, to 2.5 cm long, upto 6 partial inflorescences, heavy with profuse fruiting, pendulous; partial inflorescence 30 cm long; primary sheaths closely sheathing, completely armed with sturdy spines; rachillae to 13 cm long; strongly recurved



C. Renuka et al.

.....**i**

1

Fig. 1. *Calamus neelagiricus* Renuka: A. A portion of stem with leaf sheath; B. Portions of a leaf; C. A partial infructescence; D. Fruit; E. Seed .

.

Rheedea 7(2), 1997

A new species of Calamus (Arecaceae) from Silent Valley, Kerala, India

.

and coiled; secondary sheaths closely sheathing. Fruit oblate, 1×0.5 cm; fruiting perianth pedicellate, scales straw yellow when dry, channelled in the middle, in 21 series. Endosperm ruminate.

Etymology: The new species is named after the Neelagiri Hills from where the specimen collected.

This species comes under the group VIII of Beccari (1908). In vegetative characters this species is very close to C. gamblei and C. delessertianus, but lack cilia on the veins of the leaflets which is characteristic of the latter. From C. gamblei it can be distinguished from the shape and colour of the fruit which is oblate and straw yellow. In C. gamblei the basal part of the fruit is tapering and is golden yellow.

Acknowledgements

We wish to express our grateful thanks to Dr. K.S.S. Nair, Director, Kerala Forest Research Institute, for giving the necessary facilities, Dr. J. F. Veldkamp, Rijksherbarium, Leiden for the Latin diagnosis, and Ms. Sheeja Remani for the line drawing.

Literature cited

- Beccari, O. 1908. Asiatic palms: Lepidocaryeae. Part I. The species of Calamus. Ann. Roy. Bot. Gard. Calcutta 11: 1-518, Pl. I-II, 1-238.
- Fernandez, R. R. & A. C. Dey. 1970. A new species of *Calamus* from Western Ghats. *Indian For.* 96: 223-225.

Fischer, C. E. C. 1931. Palmae. In: J.S. Gamble, Flora of the Presidency of Madras. London. 1: 1553-1568.

Renuka, C. 1992. Rattans of the Western Ghats - A taxonomic manual. KFRI, Kerala. India.